

# Axillary lymph node and breast metastasis from early stage endometrial adenocarcinoma after a 4-year period: A case report and a retrospective study

huang chunyu<sup>1</sup>, chen liwen<sup>1</sup>, huang donglan<sup>1</sup>, wang hongmei<sup>1</sup>, zhang weijun<sup>1</sup>, shi jianjun<sup>1</sup>, and yuan yawei<sup>1</sup>

<sup>1</sup>Guangzhou Medical University Affiliated Cancer Hospital


October 26, 2022



## Abstract


Background. Endometrial adenocarcinoma (EC) with early stage, high grade is rare seen metastasis. Here, we describe a unique case of axillary lymph node and breast metastasis from high grade endometrial adenocarcinoma after a 4-year period. Cases. We presented a case of a 60-year-old woman with EC and received surgery in 2013. However, a metastatic tumor appeared at the axillary lymph node and the breast 4 years after surgery. Then she received an operation and 6 cycles adjuvant chemotherapy. 31 months following, she is alive without disease. Conclusions. A correct diagnosis should be made by combining the patient's past clinical history, morphologic and immunohistochemical test results. Knowledge of this rare metastatic site will help the physician make the correct diagnosis.

## Hosted file

manuscript.doc available at <https://authorea.com/users/517792/articles/592201-axillary-lymph-node-and-breast-metastasis-from-early-stage-endometrial-adenocarcinoma-after-a-4-year-period-a-case-report-and-a-retrospective-study>







Hosted file

Table 1.doc available at <https://authorea.com/users/517792/articles/592201-axillary-lymph-node-and-breast-metastasis-from-early-stage-endometrial-adenocarcinoma-after-a-4-year-period-a-case-report-and-a-retrospective-study>

Hosted file

Table 2.docx available at <https://authorea.com/users/517792/articles/592201-axillary-lymph-node-and-breast-metastasis-from-early-stage-endometrial-adenocarcinoma-after-a-4-year-period-a-case-report-and-a-retrospective-study>