

# Molecular ecology and microbiomes in the wild: methodological advances, common pitfalls and future directions

Nick Fountain-Jones<sup>1</sup>, Tatiana Giraud<sup>2</sup>, Lucie Zinger<sup>3</sup>, Holly Bik<sup>4</sup>, Simon Creer<sup>5</sup>, and Elin Videvall<sup>6</sup>

<sup>1</sup>University of Tasmania

<sup>2</sup>Université Paris-Sud

<sup>3</sup>Institut de Biologie de l'ENS (IBENS), Département de Biologie, École normale supérieure, CNRS, INSERM, Université PSL, Paris, France

<sup>4</sup>University of Georgia

<sup>5</sup>Bangor University

<sup>6</sup>Brown University

September 27, 2023

## Abstract

The study of microbiomes across organisms and environments has become a prominent focus in molecular ecology. This perspective article explores methodological advancements, common challenges and future directions in the field. Key research areas include understanding the drivers of microbiome community assembly, linking microbiome composition to host genetics, exploring microbial functions, transience, and spatial partitioning, and disentangling non-bacterial components of the microbiome. Methodological advancements, such as quantifying absolute abundances, sequencing complete genomes, and utilizing novel statistical approaches, are also useful tools for understanding complex microbial diversity patterns. Our aims are to encourage robust practices in microbiome studies and inspire researchers to explore the next frontier of this rapidly changing field.

## Introduction

The study of microbial communities across host species and environments (hereafter the ‘microbiome’) is a major focus of research in the field of molecular ecology. As such, *Molecular Ecology* and *Molecular Ecology Resources* regularly publish papers and special issues in the field. *Molecular Ecology* is at the forefront of examining the importance of microbiomes across ecosystems, from increasing our understanding of host-pathogen (e.g., Bergner et al., 2020; Wille et al., 2018) and host-symbiont interactions (e.g., Rubin et al., 2019) to investigating the impact of climate and other environmental factors on microbial populations (e.g., Santos-Júnior et al., 2022; Wu et al., 2022). Complementing these efforts, *Molecular Ecology Resources* publishes significant methodological advances that continue to shape the field (e.g., Harrison et al., 2020; Schnell et al., 2015; Stothart et al., 2021). Based on our collective experience as subject editors at *Molecular Ecology* and *Molecular Ecology Resources*, here we discuss some of the best practices and advances for analysing microbiomes, from study design to data analysis, and highlight anticipated future directions in the field. With this article we hope to inspire and encourage researchers to obtain more robust insights from microbiome data, which will enable the field to advance and tackle the new horizons enabled by recent advances in technology.

## Study design

As in any study, the sampling or experimental design of microbiome studies should include sufficient independent replicates, avoiding confounding effects as much as possible, with the samples representing appropriate ecological scales given the processes investigated. Microbiome sampling design must also be well-planned and appropriate to the specific hypothesis that is being tested. When testing hypotheses pertaining to the impact of outlier external drivers (e.g., fire, pollution events, natural disasters), studies would ideally feature samples that were collected both before and after the event, that are not confounded by habitat type, geography or physicochemistry. Before embarking on microbiome studies in the wild, particularly those of which are opportunistic (i.e., with samples originally collected for other purposes), researchers should carefully consider if autocorrelation of factors beyond their control could impede the interpretation of results. In other words, researchers must be realistic about what can be accomplished with limited sample sets, since rigorous hypothesis testing requires equally rigorous sampling protocols and study design.

In addition, the sampling of microbial communities should take into account their high heterogeneity at small spatial scales due to micro/mesoscale heterogeneity of their environment (Vos et al., 2013; Zhang et al., 2014) or neutral assembly dynamics (Woodcock et al., 2007)). For example, composite samples (i.e. pooled individual samples) can be combined prior to homogenisation and sub-sampling, in order to reduce the local, micro-scale heterogeneity if it is irrelevant to the questions being studied (George et al., 2019). Here, knowledge of how, and at what scale, the target community responds to external drivers will inform adequate sampling design. For example, a composite 0.2mg sediment sample is likely to be representative of the bacterial, archaeal, and microbial eukaryotic biospheres, but will not sample microscopic invertebrate taxa effectively, due to issues of scale (Nascimento et al., 2018). Smaller samples will contain some microscopic taxa and trace environmental DNA but they are inadequate at representing the underlying meio- and macrofaunal communities. As the target organisms grow in size, the sample volume and spatial extent of the studied area should be correspondingly expanded.

### Sample preparation and bioinformatics

In addition to the above considerations, the study design needs to account for the sensitivity and error-prone nature of many molecular-based approaches. Both shotgun metagenomics and DNA metabarcoding (i.e., amplicon sequencing of marker genes). include numerous opportunities for introducing false negatives and positives during the data generation process, starting from sample collection to the laboratory, sequencing, bioinformatics, and data analyses. Details of these issues are already largely covered in another *Molecular Ecology* editorial (Zinger et al., 2019), but to summarize briefly, some possible pitfalls include sample contamination stemming from the field or lab environment (de Goffau et al., 2018; Salter et al., 2014), extraction/PCR amplification biases, and errors generated during PCR and sequencing. Technical considerations, such as sample volume and choice of lab reagents, are, in many cases, the result of a compromise between the research question, logistical feasibility, time and available funds (Taberlet et al., 2018). However, any compromise of the protocol should still allow one to appropriately address the research question. In addition, we want to re-emphasize the importance of adequately describing the whole data production workflow in the methods section of manuscripts (e.g., primer sequences, polymerase, molecular labelling strategy). Environmental or lab contamination is a particularly large problem for samples with low microbial biomass (Eisenhofer et al., 2019), and the collection of such low biomass samples cannot be avoided in many study designs (e.g. host-associated microbiomes of small organisms, or depauperate environmental habitats). The sequencing of negative controls (and potentially also positive controls and technical replicates) alongside experimental samples is important for quantifying errors and artefacts (e.g., Davis et al., 2018), and can improve data curation procedures through tuned, experiment-specific criteria, including for samples with low microbial biomass. While there is more than one way to implement such efforts, a thorough description of the controls, a rationale for including them, and the ways they are integrated into data analysis, are essential practices of good microbiome science (Hakimzadeh et al., 2023).

One overlooked problem in microbiome studies is cross-contamination between samples during library preparation procedures (Kim et al., 2017; Zinger et al., 2019), which can result in an artificial reduction in beta diversity (i.e., compositional differences between samples) and an increase in alpha diversity. Such

cross-contamination can occur during the PCR plate preparation process through pipetting errors or aerosol production. Random positioning of samples in the PCR plates provides a relatively simple approach to reduce this problem (Minich et al., 2019; Taberlet et al., 2018). More often, - and insidiously - cross-contamination can occur during the PCR cycles, a bias referred to as tag-switches (Carlsen et al., 2012; Esling et al., 2015), tag-jumps (Schnell et al., 2015), or, more recently in the microbiome literature, cross-talks (Edgar, 2018; Minich et al., 2019). During this laboratory step, amplicon molecules from different samples can recombine within conserved primer sequences, resulting in the production of new molecules containing the genuine DNA sequence, but the wrong sample/barcode label. As a consequence, the most abundant taxa will be detected in many samples, including the negative controls (Esling et al., 2015; Minich et al., 2019; Taberlet et al., 2018), preventing simple removal of all taxa occurring in controls as a fix for field and lab contamination. Alleviating the problem of tag-switches can be achieved with modified library preparation protocols (e.g., Carøe & Bohmann, 2020), appropriate sample labelling strategies and *a posteriori* using the information contained in both samples and negative controls (Bohmann et al., 2022; Hakimzadeh et al., 2023).

One limitation in microbiome studies, using either DNA metabarcoding or metagenomics, is the compositional nature of the sequence data (i.e., described as proportions or probabilities, not absolute number of molecules). Like other count-based sequencing approaches, this limitation arises because the concentration of PCR products is standardised prior to sequencing and the number of molecules read is limited by the sequencing platform (Gloor et al., 2017). As a result, we are unable to obtain data on absolute abundances and biomass for the different microbial members of the community. However, exciting new developments are emerging to overcome these limitations, relying on known reference values of DNA molecule abundances, allowing simple conversion of relative abundance into absolute values. Two broad classes of methods based on this approach have emerged: (i) quantification of target markers using q/ddPCR prior to metabarcoding (Barlow et al., 2020; Callahan et al., 2019; Ji et al., 2019) and (ii) introduction of exogenous DNA molecules spike-ins (i.e. DNA molecules of known sequence and quantity to calibrate measurements). Related to the latter, a host-associated microbiome PCR approach (HamPCR, Lundberg et al., 2021) represents a promising method to assess the ratio of the microbial population size relative to the amount of host tissue (i.e. microbial load).

Despite obtaining a better estimate of the absolute number of molecules in a sample, it is still challenging to convert this number into the actual number of microbial cells. The calculation is often difficult because most gold-standard barcoding genes used for bacteria, fungi, and protists have multiple copies in the genome, with precise numbers varying across taxa and in unpredictable ways (Louca et al., 2018). Another problem is that some of the retrieved molecules can be derived from extracellular DNA or DNA adsorbed on cell debris or particles, i.e. correspond to non-living organisms (Torti et al., 2015). The proportion of extracellular DNA in the environment is often not known but can be estimated with different approaches (reviewed by Nagler et al., 2022). Further, because extracellular DNA is often degraded, approaches including long-read sequencing targeting larger genomic regions will likely overcome this issue. Approaches that are able to quantify or eliminate extracellular DNA can prove useful when having correct snapshots of the microbial community is crucial (e.g., when studying short-term processes with repeated observations capturing microbiome variation within host individuals) but are likely less relevant when studying processes operating at larger temporal scales (e.g., microbiome response to climate change).

Lastly, incorporating site-occupancy modelling in microbiome studies presents an exciting avenue to quantify measurement uncertainty and to account for imperfect detection (e.g., Ficetola et al., 2015; McClenaghan et al., 2020; Willoughby et al., 2016). Site-occupancy models use data collected over multiple visits to sites (or across multiple technical/biological replicates) to quantify how likely it is to detect a taxon when it is present. For microbiome studies, including both biological and technical PCR replicates can enable rigorous statistical predictions regarding the true or false positive detection of microbial species within the community. Further, these predictions can be utilised to improve study design (Fukaya et al., 2022; McClenaghan et al., 2020). How many replicates of each type is required is an open question, although biological replicates may improve detection probabilities (Willoughby et al., 2016).


## Beyond estimating diversity: Exciting advances in statistics

Modelling advances in community ecology offer exciting opportunities to understand the complex patterns in microbial diversity and complement robust sampling designs (e.g., Grantham et al., 2020; Trego et al., 2022). In addition, novel methods for analyzing amplicon sequencing data are continuously emerging, primarily focused on the human gut microbiome but adaptable to other microbial ecology fields with suitable study designs and datasets (e.g., Trego et al., 2022). Broadly, these tools can be categorized into quantifying community assembly processes, mapping occurrence networks, capturing spatial/temporal dynamics, integrating multi-omics, identifying differentially abundant taxa, finding species-environment associations, and predicting functional patterns (Trego et al., 2022). However, despite the frequent use of high-throughput sequencing, there has been a slow uptake of these new analytical techniques, and many studies do not go much beyond basic comparisons of alpha and beta diversity estimates across samples. While important inferences can be made by examining overall patterns of composition and diversity (e.g., Grosser et al., 2019; Motta et al., 2018), they offer only a starting point toward having a more mechanistic understanding of the ecological drivers of microbiome variation (Shade, 2017).

Common analytical approaches to quantify differences in beta diversity across microbiome samples, such as the permutational multivariate analysis of variance (PERMANOVA), are algorithmic (i.e., not based on a statistical model) and do not explicitly account for uncertainty in ecological data (Björk et al., 2018; Warton et al., 2012, 2015). Importantly, making inferences about microbiome variation is often difficult using algorithmic distance-based approaches (Björk et al., 2018; Warton et al., 2012). Model-based approaches such as joint species distribution models (JSDMs) or stacked models (Powell-Romero et al., 2023) are multi-response extensions of generalized linear mixed models (GLMMs) that can overcome some of the limitations of the algorithmic methods to elucidate patterns of microbiome variation (e.g., Björk et al., 2018; Grantham et al., 2020). Often using a Bayesian framework, JSDMs simultaneously analyze multiple species and environmental variables, allowing for the assessment of community-level responses to environmental change and host effects (Björk et al., 2018; Ovaskainen et al., 2017; Pollock et al., 2014). JSDMs can i) incorporate information on species traits and phylogenetic relatedness, improving estimation accuracy and power when there is a phylogenetic signal (Ovaskainen et al., 2017), and ii) analyse patterns of taxon covariance to infer microbial co-occurrence networks (Björk et al., 2018; Fountain-Jones et al., 2020, 2023). Microbial co-occurrence networks are valuable tools in microbiome science, as they offer insights (but see *Current gaps and future directions* below) into the associations among microbial taxa, enhancing our understanding of microbial community dynamics and functioning. JSDM-based co-occurrence networks have an added advantage of interpretation as the major environmental and host effects shaping microbial presences are controlled for (i.e., an inferred association between microbes is then not likely a mere product of a shared environmental response). However, GLMM-based JSDM co-occurrence networks cannot untangle the relative roles of taxa associations, and environmental or host effects (Clark et al., 2018; Fountain-Jones et al., 2020) and tend to not scale well with large datasets (Pichler & Hartig, 2021). Approaches such as conditional random fields (CRF, Clark et al., 2018), multi-response interpretable machine learning (mrIML, Fountain-Jones et al., 2021), MIMIX (Microbiome MIXed Model, Grantham et al., 2020) and scalable JSDMs (sjSDM, Pichler & Hartig, 2021) can overcome these limitations. Importantly, approaches such as MrIML and MiMiX allow for predictions and treatment effects to be extracted for individual taxa, which can be useful if researchers have a set of focal taxa. We note that these methods are not appropriate in all situations. For particularly large datasets (thousands of samples), new distance-based methods such as D-MANOVA (Chen & Zhang, 2021) or multivariate distance matrix regression (MDMR, Zapala & Schork, 2012) may be better options. Boshuizen & te Beest (2023) have provided a complete guide of the pitfalls in analysing amplicon data. While the tools mentioned here represent only a tiny fraction of the potential methods available, we encourage readers to go beyond diversity metrics and differentially abundant taxa to gain more mechanistic insights into microbiome data from wild species.

Incorporating some of the methodological advances in bioinformatics and statistics, coupled with robust study design, and rigorous laboratory techniques, will improve current research efforts in the field (see Fig. 1 for a summary). Moreover, taking into consideration both the limitations and opportunities of these various

approaches allow us to open up new exciting avenues in the field of microbiome ecology research.



**Fig. 1:** Overview of some of the considerations and techniques that can be employed across a general microbiome workflow. JSDMs: Joint species distribution models. q/dd PCR quantitative/ double digest PCR.

### Current gaps and future directions in microbiome ecology research

Given the above considerations, how do we move the field forward? Here we outline some applicable research directions that will generate significant impact, by helping to close some of the most pressing knowledge gaps in the near future.

**1. Obtaining a better understanding of the ecological, evolutionary and mechanistic drivers of microbiome community assembly.** A key research gap in microbiome ecology is the need for a comprehensive understanding of the drivers of community assembly. While significant progress has been made regarding microbiomes associated with humans and model organisms (e.g., *Drosophila* or *Arabidopsis*), further study on non-model animal and plant systems is required. Included within this goal is the investigation of phylosymbiosis – the topological congruence between host phylogenetic distance and the compositional similarity patterns of their associated microbiota (Brown et al., 2023); a pattern that can arise from both ecological and evolutionary processes. Mechanistic studies elucidating the specific processes that govern microbial transmission, colonization, competition and succession will help explain the presence or absence of phylosymbiosis signals across host species or populations, and are more broadly essential for a deeper understanding of microbiome assembly (Coyte et al., 2021).

**2. Linking microbiome composition and host genetics.** To gain insights into the ability of hosts to select specific microbes that may benefit their health or reproduction, future research should aim to link host-associated microbiome composition and diversity with genes and genomic regions of hosts. This hologenomic approach will enable the identification of host genetic factors, such as immune genes, that are key players in shaping the host microbiome and ultimately the resulting host phenotypes. By integrating both host genome and microbiome data, researchers can make progress at uncovering host-microbiome interactions (Sutherland et al., 2022; West et al., 2023). Incorporation of long-read sequencing and hybrid assembly approaches which utilize both short and long reads now offer exciting opportunities for advancing this research area.

**3. Linking microbiomes to host traits and phenotypes.** Understanding the connection between host-associated microbiomes and host phenotypes is a related and important research avenue. Investigating the influence of the microbiome on host development,

behavior, metabolism, and life-history traits can therefore provide valuable insights (e.g., Bestion et al., 2017; Wood et al., 2022). For example, some nematode species are known to demonstrate extreme phenotypic plasticity in response to environmental cues (e.g. chemical or bacterial stimuli (Hauquier et al., 2017; Sommer et al., 2017), and one open question is whether microbial taxa play an integral role in initiating such host developmental switch genes (which are themselves under epigenetic control in the case of *Pristionchus* spp. fig nematodes). By integrating microbiota data with detailed trait measurements of diverse hosts, researchers will be able to identify host phenotypes associated with certain microbiome compositions and start to unravel the underlying mechanisms of how certain microbes can influence the phenotypes of hosts, and vice versa.

**4. Exploring microbial functions within host-associated microbiomes.** While microbial community composition and diversity have been extensively studied in microbiome ecology, there is a significant need to explore the functional attributes of whole communities, localized populations, and individual microorganisms (genes and pangenomes) within a microbiome. Investigating microbial functions, such as metabolic pathways and molecular interactions between members of the microbiome and with the host, can provide important insights into the contributions of specific microbial taxa/consortia and their functional roles in host and ecosystem health (Béchade et al., 2023; e.g., Hicks et al., 2018; Karmacharya et al., 2019). Furthermore, isolation and culturing of microbial strains can provide complementary information not otherwise accessible through community -omics alone (e.g., physiological profiling of microbial growth rates and chemical/antibiotic sensitivity), while also paving the way for future experimental work using such host-associated microbial isolates.

**5. Disentangle the role of the non-bacterial components of the microbiome.** Although bacteria have been the primary focus of microbiome ecology research due to their overwhelming abundance, other components such as viruses, fungi, or protists play crucial roles in host-microbe interactions and ecosystem functioning (Jervis et al., 2021; Raghwani et al., 2023). Future investigations that include these non-bacterial components will allow us to more comprehensively understand the dynamics of the microbiome as a whole community, its interaction with hosts, and its role in the ecosystem.

**6. Elucidate the role of host-associated microbiomes in wildlife disease and conservation ecology.** Understanding the role of microbiomes in biological conservation, such as wildlife disease susceptibility and resistance, is an emerging and timely field of research within microbiome ecology. Investigating the interactions between host genetics, environmental factors and microbial communities can shed light on disease dynamics and the mechanisms through which microbiomes modulate host immune responses in wildlife populations (e.g., Bozzi et al., 2021; Gao et al., 2021; Jervis et al., 2021). Likewise, studies linking environmental microbiomes with land use, habitat fragmentation, and climate change can provide important information on how to address ecosystem challenges in a changing world.

**7. Disentangling diet-microbiome associations.** Diet is a well-known driver of microbiome composition in hosts, but the mechanisms by which components of the diet promote certain taxa and ultimately influences host health remain unclear. Therefore, the complexity of diet-microbiome associations requires further investigation (Kartzinel et al., 2019). Future research should aim to unravel the specific dietary components that shape microbial communities, how diet diversity is related to microbial diversity, and the mechanisms through which these interactions ultimately influence host health. Longitudinal studies and controlled dietary interventions can provide valuable insights into the specifics of diet-microbiome relationships (Couch et al., 2021).

**8. Unravelling microbial interactions within the microbiome.** Elucidating the nature of microbial interactions is crucial to understand the dynamics of microbial ecosystems. Patterns of co-occurrence are often used to evaluate microbial interactions (e.g., competition), yet doing so is problematic (see Blanchet et al. (2020)). Interactions are highly scale-dependent, which poses unique challenges for microbial communities with fine spatial structuring (Goberna & Verdú, 2022; Peng et al., 2023). Future experimental and observational studies at relevant scales, with large numbers of samples across time and including robust measures of abundance, will be able to better quantify microbial interactions (Blanchet et al., 2020). Statistical advances utilising generalized Lotka–Volterra models across time-series (Stein et al., 2013), or employing conditional probabilities to more directly capture how taxa relate to each other will also help infer interactions (Blanchet et al., 2020).

**9. Determine microbial strain diversity and evolution within hosts.** Microbial communities are often highly heterogeneous and different strains of a single microbial species can exhibit significant genetic and functional variability (Anderson & Bisanz, 2023; Goyal et al., 2022). Investigating microbial genetic diversity and the role of horizontal gene

transfer are therefore crucial to better understand the adaptive processes and functional implications within a microbiome (Barreto & Gordo, 2023). For example, a shift in *Escherichia coli* clones was documented in the gut microbiome of ageing mice, and these were characterized by an increase in bacterial mutations targeting stress-response genes (Barreto et al., 2020). Integrating high-resolution genomic techniques with longitudinal and repeated sampling schemes can capture key patterns in temporal variation among microbial communities and significantly improve our understanding of how microbes evolve within individual hosts or specific environments.

## Conclusions

We are currently at a major turning point on how we can derive valuable insights on the ecological processes shaping non-model and environmental microbiomes in the wild. The adoption of robust laboratory and bioinformatic techniques, together with sophisticated statistical approaches, enhance our ability to gain deeper insights into the factors influencing microbiome variation and the intricate relationships among microbial taxa. However, all of these techniques rely on robust study designs and appropriate sampling scales to address specific research questions. Ultimately, this perspective piece serves as a broad outline of some of the considerations that ought to be considered in the field of microbiome ecology. Hopefully, with robust practices, we can, in turn, start to untangle the complex processes acting on these incredibly important but understudied communities.

## Acknowledgments

We would like to thank Jacob A. Russell, Eric Coissac and Kayla King for their useful discussions and edits to the text.

## Author Contributions

NFJ and EV conceived the project. All authors contributed to writing the first draft of the manuscript and subsequent editing.

## Conflict of interest

None.

## References

- Anderson, B. D., & Bisanz, J. E. (2023). Challenges and opportunities of strain diversity in gut microbiome research. *Frontiers in Microbiology* , 14 . <https://www.frontiersin.org/articles/10.3389/fmicb.2023.1117122>
- Barlow, J. T., Bogatyrev, S. R., & Ismagilov, R. F. (2020). A quantitative sequencing framework for absolute abundance measurements of mucosal and luminal microbial communities. *Nature Communications* , 11 (1), Article 1. <https://doi.org/10.1038/s41467-020-16224-6>
- Barreto, H. C., & Gordo, I. (2023). Intrahost evolution of the gut microbiota. *Nature Reviews Microbiology* , 21 (9), Article 9. <https://doi.org/10.1038/s41579-023-00890-6>
- Barreto, H. C., Sousa, A., & Gordo, I. (2020). The Landscape of Adaptive Evolution of a Gut Commensal Bacteria in Aging Mice. *Current Biology* , 30 (6), 1102-1109.e5. <https://doi.org/10.1016/j.cub.2020.01.037>
- Béchade, B., Cabuslay, C. S., Hu, Y., Mendonca, C. M., Hassanpour, B., Lin, J. Y., Su, Y., Fiers, V. J., Anandarajan, D., Lu, R., Olson, C. J., Duplais, C., Rosen, G. L., Moreau, C. S., Aristilde, L., Wertz, J. T., & Russell, J. A. (2023). Physiological and evolutionary contexts of a new symbiotic species from the nitrogen-recycling gut community of turtle ants. *The ISME Journal* , 17 (10), 1751–1764. <https://doi.org/10.1038/s41396-023-01490-1>
- Bergner, L. M., Orton, R. J., Benavides, J. A., Becker, D. J., Tello, C., Biek, R., & Streicker, D. G. (2020). Demographic and environmental drivers of metagenomic viral diversity in vampire bats. *Molecular Ecology* , 29 (1), 26–39. <https://doi.org/10.1111/mec.15250>

- Bestion, E., Jacob, S., Zinger, L., Di Gesu, L., Richard, M., White, J., & Cote, J. (2017). Climate warming reduces gut microbiota diversity in a vertebrate ectotherm. *Nature Ecology & Evolution* , 1 (6), 161. <https://doi.org/10.1038/s41559-017-0161>
- Björk, J. R., Hui, F. K. C., O’Hara, R. B., & Montoya, J. M. (2018). Uncovering the drivers of host-associated microbiota with joint species distribution modelling. *Molecular Ecology* , 27 (12), 2714–2724. <https://doi.org/10.1111/mec.14718>
- Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters* , 23 (7), 1050–1063. <https://doi.org/10.1111/ele.13525>
- Bohmann, K., Elbrecht, V., Carøe, C., Bista, I., Leese, F., Bunce, M., Yu, D. W., Seymour, M., Dumbrell, A. J., & Creer, S. (2022). Strategies for sample labelling and library preparation in DNA metabarcoding studies. *Molecular Ecology Resources* , 22 (4), 1231–1246. <https://doi.org/10.1111/1755-0998.13512>
- Boshuizen, H. C., & te Beest, D. E. (2023). Pitfalls in the statistical analysis of microbiome amplicon sequencing data. *Molecular Ecology Resources* , 23 (3), 539–548. <https://doi.org/10.1111/1755-0998.13730>
- Bozzi, D., Rasmussen, J. A., Carøe, C., Sveier, H., Nordøy, K., Gilbert, M. T. P., & Limborg, M. T. (2021). Salmon gut microbiota correlates with disease infection status: Potential for monitoring health in farmed animals. *Animal Microbiome* , 3 (1), 30. <https://doi.org/10.1186/s42523-021-00096-2>
- Brown, B. R. P., Goheen, J. R., Newsome, S. D., Pringle, R. M., Palmer, T. M., Khasoha, L. M., & Kartzinel, T. R. (2023). Host phylogeny and functional traits differentiate gut microbiomes in a diverse natural community of small mammals. *Molecular Ecology* , 32 (9), 2320–2334. <https://doi.org/10.1111/mec.16874>
- Callahan, B. J., Wong, J., Heiner, C., Oh, S., Theriot, C. M., Gulati, A. S., McGill, S. K., & Dougherty, M. K. (2019). High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution. *Nucleic Acids Research* , 47 (18), e103. <https://doi.org/10.1093/nar/gkz569>
- Carlsen, T., Aas, A. B., Lindner, D., Vrålstad, T., Schumacher, T., & Kauserud, H. (2012). Don’t make a mista(g)ke: Is tag switching an overlooked source of error in amplicon pyrosequencing studies? *Fungal Ecology* , 5 (6), 747–749. <https://doi.org/10.1016/j.funeco.2012.06.003>
- Carøe, C., & Bohmann, K. (2020). Tagsteady: A metabarcoding library preparation protocol to avoid false assignment of sequences to samples. *Molecular Ecology Resources* , 20 (6), 1620–1631. <https://doi.org/10.1111/1755-0998.13227>
- Chen, J., & Zhang, X. (2021). D-MANOVA: Fast distance-based multivariate analysis of variance for large-scale microbiome association studies. *Bioinformatics* , 38 (1), 286–288. <https://doi.org/10.1093/bioinformatics/btab498>
- Clark, N. J., Wells, K., & Lindberg, O. (2018). Unravelling changing interspecific interactions across environmental gradients using Markov random fields. *Ecology* , 99 (6), 1277–1283. <https://doi.org/10.1002/ecy.2221>
- Couch, C. E., Stagaman, K., Spaan, R. S., Combrink, H. J., Sharpton, T. J., Beechler, B. R., & Jolles, A. E. (2021). Diet and gut microbiome enterotype are associated at the population level in African buffalo. *Nature Communications* , 12 (1), Article 1. <https://doi.org/10.1038/s41467-021-22510-8>
- Coyte, K. Z., Rao, C., Rakoff-Nahoum, S., & Foster, K. R. (2021). Ecological rules for the assembly of microbiome communities. *PLOS Biology* , 19 (2), e3001116. <https://doi.org/10.1371/journal.pbio.3001116>
- Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018). Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome* , 6 (1), 226. <https://doi.org/10.1186/s40168-018-0605-2>
- de Goffau, M. C., Lager, S., Salter, S. J., Wagner, J., Kronbichler, A., Charnock-Jones, D. S., Peacock, S. J., Smith, G. C. S., & Parkhill, J. (2018). Recognizing the reagent microbiome. *Nature Microbiology* , 3 (8), Article 8. <https://doi.org/10.1038/s41564-018-0202-y>

- Edgar, R. C. (2018). *UNCROSS2: Identification of cross-talk in 16S rRNA OTU tables* (p. 400762). bioRxiv. <https://doi.org/10.1101/400762>
- Eisenhofer, R., Minich, J. J., Marotz, C., Cooper, A., Knight, R., & Weyrich, L. S. (2019). Contamination in Low Microbial Biomass Microbiome Studies: Issues and Recommendations. *Trends in Microbiology* , *27* (2), 105–117. <https://doi.org/10.1016/j.tim.2018.11.003>
- Esling, P., Lejzerowicz, F., & Pawlowski, J. (2015). Accurate multiplexing and filtering for high-throughput amplicon-sequencing. *Nucleic Acids Research* , *43* (5), 2513–2524. <https://doi.org/10.1093/nar/gkv107>
- Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguët-Covex, C., De Barba, M., Gielly, L., Lopes, C. M., Boyer, F., Pompanon, F., Rayé, G., & Taberlet, P. (2015). Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources* , *15* (3), 543–556. <https://doi.org/10.1111/1755-0998.12338>
- Fountain-Jones, N., Kozakiewicz, C., Forester, B., Landguth, E., Carver, S., Charleston, M., Gagne, R., Greenwell, B., Kraberger, S., Trumbo, D., Mayer, M., Clark, N., & Machado, G. (2021). MrIML: Multi-response interpretable machine learning to map genomic landscapes. *Molecular Ecology Resources* , *21* , 2766–2781. <https://doi.org/10.22541/au.160855820.09604024/v1>
- Fountain-Jones, N. M., Clark, N. J., Kinsley, A. C., Carstensen, M., Forester, J., Johnson, T. J., Miller, E. A., Moore, S., Wolf, T. M., & Craft, M. E. (2020). Microbial associations and spatial proximity predict North American moose (*Alces alces*) gastrointestinal community composition. *Journal of Animal Ecology* , *89* (3). <https://doi.org/10.1111/1365-2656.13154>
- Fountain-Jones, N. M., Khoo, B. S., Rau, A., Berman, J. D., Burton, E. N., & Oliver, J. D. (2023). Positive associations matter: Microbial relationships drive tick microbiome composition. *Molecular Ecology* , *32* (14), 4078–4092. <https://doi.org/10.1111/mec.16985>
- Fukaya, K., Kondo, N. I., Matsuzaki, S. S., & Kadoya, T. (2022). Multispecies site occupancy modelling and study design for spatially replicated environmental DNA metabarcoding. *Methods in Ecology and Evolution* , *13* (1), 183–193. <https://doi.org/10.1111/2041-210X.13732>
- Gao, M., Xiong, C., Gao, C., Tsui, C. K. M., Wang, M.-M., Zhou, X., Zhang, A.-M., & Cai, L. (2021). Disease-induced changes in plant microbiome assembly and functional adaptation. *Microbiome* , *9* (1), 187. <https://doi.org/10.1186/s40168-021-01138-2>
- George, P. B. L., Lallias, D., Creer, S., Seaton, F. M., Kenny, J. G., Eccles, R. M., Griffiths, R. I., Lebron, I., Emmett, B. A., Robinson, D. A., & Jones, D. L. (2019). Divergent national-scale trends of microbial and animal biodiversity revealed across diverse temperate soil ecosystems. *Nature Communications* , *10* (1), 1107. <https://doi.org/10.1038/s41467-019-09031-1>
- Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., & Egozcue, J. J. (2017). Microbiome Datasets Are Compositional: And This Is Not Optional. *Frontiers in Microbiology* , *8* . <https://www.frontiersin.org/articles/10.3389/fmicb.2017.02224>
- Goberna, M., & Verdú, M. (2022). Cautionary notes on the use of co-occurrence networks in soil ecology. *Soil Biology and Biochemistry* , *166* , 108534. <https://doi.org/10.1016/j.soilbio.2021.108534>
- Goyal, A., Bittleston, L. S., Leventhal, G. E., Lu, L., & Cordero, O. X. (2022). Interactions between strains govern the eco-evolutionary dynamics of microbial communities. *ELife* , *11* , e74987. <https://doi.org/10.7554/eLife.74987>
- Grantham, N. S., Guan, Y., Reich, B. J., Borer, E. T., & Gross, K. (2020). MIMIX: A Bayesian Mixed-Effects Model for Microbiome Data From Designed Experiments. *Journal of the American Statistical Association* , *115* (530), 599–609. <https://doi.org/10.1080/01621459.2019.1626242>

- Grosser, S., Sauer, J., Paijmans, A. J., Caspers, B. A., Forcada, J., Wolf, J. B. W., & Hoffman, J. I. (2019). Fur seal microbiota are shaped by the social and physical environment, show mother–offspring similarities and are associated with host genetic quality. *Molecular Ecology* , 28 (9), 2406–2422. <https://doi.org/10.1111/mec.15070>
- Hakimzadeh, A., Abdala Asbun, A., Albanese, D., Bernard, M., Buchner, D., Callahan, B., Caporaso, J. G., Curd, E., Djemiel, C., Brandström Durling, M., Elbrecht, V., Gold, Z., Gweon, H. S., Hajibabaei, M., Hildebrand, F., Mikryukov, V., Normandeau, E., Özkurt, E., M. Palmer, J., . . . Anslan, S. (n.d.). A pile of pipelines: An overview of the bioinformatics software for metabarcoding data analyses. *Molecular Ecology Resources* , n/a (n/a). <https://doi.org/10.1111/1755-0998.13847>
- Harrison, J. G., John Calder, W., Shuman, B., & Alex Buerkle, C. (2020). The quest for absolute abundance: The use of internal standards for DNA-based community ecology. *Molecular Ecology Resources* , 1755-0998.13247. <https://doi.org/10.1111/1755-0998.13247>
- Hauquier, F., Leliaert, F., Rigaux, A., Derycke, S., & Vanreusel, A. (2017). Distinct genetic differentiation and species diversification within two marine nematodes with different habitat preference in Antarctic sediments. *BMC Evolutionary Biology* , 17 (1), 120. <https://doi.org/10.1186/s12862-017-0968-1>
- Hicks, A. L., Lee, K. J., Couto-Rodriguez, M., Patel, J., Sinha, R., Guo, C., Olson, S. H., Seimon, A., Seimon, T. A., Ondzie, A. U., Karesh, W. B., Reed, P., Cameron, K. N., Lipkin, W. I., & Williams, B. L. (2018). Gut microbiomes of wild great apes fluctuate seasonally in response to diet. *Nature Communications* , 9 , 1786. <https://doi.org/10.1038/s41467-018-04204-w>
- Jervis, P., Pintanel, P., Hopkins, K., Wierzbicki, C., Shelton, J. M. G., Skelly, E., Rosa, G. M., Almeida-Reinoso, D., Eugenia-Ordóñez, M., Ron, S., Harrison, X., Merino-Viteri, A., & Fisher, M. C. (2021). Post-epizootic microbiome associations across communities of neotropical amphibians. *Molecular Ecology* , 30 (5), 1322–1335. <https://doi.org/10.1111/mec.15789>
- Ji, B. W., Sheth, R. U., Dixit, P. D., Huang, Y., Kaufman, A., Wang, H. H., & Vitkup, D. (2019). Quantifying spatiotemporal variability and noise in absolute microbiota abundances using replicate sampling. *Nature Methods* , 16 (8), Article 8. <https://doi.org/10.1038/s41592-019-0467-y>
- Karmacharya, D., Manandhar, P., Manandhar, S., Sherchan, A. M., Sharma, A. N., Joshi, J., Bista, M., Bajracharya, S., Awasthi, N. P., Sharma, N., Llewellyn, B., Waits, L. P., Thapa, K., Kelly, M. J., Vuyisich, M., Starckenburg, S. R., Hero, J.-M., Hughes, J., Wultsch, C., . . . Sinha, A. K. (2019). Gut microbiota and their putative metabolic functions in fragmented Bengal tiger population of Nepal. *PLoS ONE* , 14 (8). <https://doi.org/10.1371/journal.pone.0221868>
- Kartzinel, T. R., Hsing, J. C., Musili, P. M., Brown, B. R. P., & Pringle, R. M. (2019). Covariation of diet and gut microbiome in African megafauna. *Proceedings of the National Academy of Sciences of the United States of America* , 116 (47), 23588–23593. <https://doi.org/10.1073/pnas.1905666116>
- Kim, D., Hofstaedter, C. E., Zhao, C., Mattei, L., Tanes, C., Clarke, E., Lauder, A., Sherrill-Mix, S., Chehoud, C., Kelsen, J., Conrad, M., Collman, R. G., Baldassano, R., Bushman, F. D., & Bittinger, K. (2017). Optimizing methods and dodging pitfalls in microbiome research. *Microbiome* , 5 (1), 52. <https://doi.org/10.1186/s40168-017-0267-5>
- Louca, S., Doebeli, M., & Parfrey, L. W. (2018). Correcting for 16S rRNA gene copy numbers in microbiome surveys remains an unsolved problem. *Microbiome* , 6 (1), 41. <https://doi.org/10.1186/s40168-018-0420-9>
- Lundberg, D. S., Pramoj Na Ayutthaya, P., Strauss, A., Shirsekar, G., Lo, W.-S., Lahaye, T., & Weigel, D. (2021). Host-associated microbe PCR (hamPCR) enables convenient measurement of both microbial load and community composition. *ELife* , 10 , e66186. <https://doi.org/10.7554/eLife.66186>
- McClenaghan, B., Compson, Z. G., & Hajibabaei, M. (2020). Validating metabarcoding-based biodiversity assessments with multi-species occupancy models: A case study using coastal marine eDNA. *PLOS ONE* ,

15 (3), e0224119. <https://doi.org/10.1371/journal.pone.0224119>

Minich, J. J., Sanders, J. G., Amir, A., Humphrey, G., Gilbert, J. A., & Knight, R. (2019). Quantifying and Understanding Well-to-Well Contamination in Microbiome Research. *MSystems* , 4 (4), 10.1128/msystems.00186-19. <https://doi.org/10.1128/msystems.00186-19>

Motta, E. V. S., Raymann, K., & Moran, N. A. (2018). Glyphosate perturbs the gut microbiota of honey bees. *Proceedings of the National Academy of Sciences* , 115 (41), 10305–10310. <https://doi.org/10.1073/pnas.1803880115>

Nagler, M., Podmirseg, S. M., Ascher-Jenull, J., Sint, D., & Traugott, M. (2022). Why eDNA fractions need consideration in biomonitoring. *Molecular Ecology Resources* , 22 (7), 2458–2470. <https://doi.org/10.1111/1755-0998.13658>

Nascimento, F. J. A., Lallias, D., Bik, H. M., & Creer, S. (2018). Sample size effects on the assessment of eukaryotic diversity and community structure in aquatic sediments using high-throughput sequencing. *Scientific Reports* , 8 (1), Article 1. <https://doi.org/10.1038/s41598-018-30179-1>

Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., Roslin, T., & Abrego, N. (2017). How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecology Letters* , 20 (5), 561–576. <https://doi.org/10.1111/ele.12757>

Peng, L., Hoban, J., Joffe, J., Smith, A. H., Carpenter, M., Marcelis, T., Patel, V., Lynn-Bell, N., Oliver, K. M., & Russell, J. A. (n.d.). Cryptic community structure and metabolic interactions among the heritable facultative symbionts of the pea aphid. *Journal of Evolutionary Biology* , n/a (n/a). <https://doi.org/10.1111/jeb.14216>

Pichler, M., & Hartig, F. (2021). A new joint species distribution model for faster and more accurate inference of species associations from big community data. *Methods in Ecology and Evolution* , 12 (11), 2159–2173. <https://doi.org/10.1111/2041-210X.13687>

Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O’Hara, R. B., Parris, K. M., Vesk, P. A., & McCarthy, M. A. (2014). Understanding co-occurrence by modelling species simultaneously with a Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution* , 5 (5), 397–406. <https://doi.org/10.1111/2041-210X.12180>

Powell-Romero, F., Fountain-Jones, N. M., Norberg, A., & Clark, N. J. (2023). Improving the predictability and interpretability of co-occurrence modelling through feature-based joint species distribution ensembles. *Methods in Ecology and Evolution* , 14 (1), 146–161. <https://doi.org/10.1111/2041-210X.13915>

Raghvani, J., Faust, C. L., Francois, S., Nguyen, D., Marsh, K., Raulo, A., Hill, S. C., Parag, K. V., Simmonds, P., Knowles, S. C. L., & Pybus, O. G. (n.d.). Seasonal dynamics of the wild rodent faecal virome. *Molecular Ecology* , n/a (n/a). <https://doi.org/10.1111/mec.16778>

Rubin, B. E. R., Kautz, S., Wray, B. D., & Moreau, C. S. (2019). Dietary specialization in mutualistic acacia-ants affects relative abundance but not identity of host-associated bacteria. *Molecular Ecology* , 28 (4), 900–916. <https://doi.org/10.1111/mec.14834>

Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., Turner, P., Parkhill, J., Loman, N. J., & Walker, A. W. (2014). Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biology* , 12 , 87. <https://doi.org/10.1186/s12915-014-0087-z>

Santos-Junior, C. D., Logares, R., & Henrique-Silva, F. (2022). Microbial population genomes from the Amazon River reveal possible modulation of the organic matter degradation process in tropical freshwaters. *Molecular Ecology* , 31 (1), 206–219. <https://doi.org/10.1111/mec.16222>

Schnell, I. B., Bohmann, K., & Gilbert, M. T. P. (2015). Tag jumps illuminated—Reducing sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology Resources* , 15 (6), 1289–1303.

<https://doi.org/10.1111/1755-0998.12402>

Shade, A. (2017). Diversity is the question, not the answer. *The ISME Journal* , 11 (1), 1–6. <https://doi.org/10.1038/ismej.2016.118>

Sommer, R. J., Dardiry, M., Lenuzzi, M., Namdeo, S., Renahan, T., Sieriebriennikov, B., & Werner, M. S. (2017). The genetics of phenotypic plasticity in nematode feeding structures. *Open Biology* , 7 (3), 160332. <https://doi.org/10.1098/rsob.160332>

Stein, R. R., Bucci, V., Toussaint, N. C., Buffie, C. G., Ratsch, G., Pamer, E. G., Sander, C., & Xavier, J. B. (2013). Ecological Modeling from Time-Series Inference: Insight into Dynamics and Stability of Intestinal Microbiota. *PLOS Computational Biology* , 9 (12), e1003388. <https://doi.org/10.1371/journal.pcbi.1003388>

Stothart, M. R., Greuel, R. J., Gavriiliuc, S., Henry, A., Wilson, A. J., McLoughlin, P. D., & Poissant, J. (2021). Bacterial dispersal and drift drive microbiome diversity patterns within a population of feral hindgut fermenters. *Molecular Ecology* , 30 (2), 555–571. <https://doi.org/10.1111/mec.15747>

Sutherland, J., Bell, T., Trexler, R. V., Carlson, J. E., & Lasky, J. R. (2022). Host genomic influence on bacterial composition in the switchgrass rhizosphere. *Molecular Ecology* , 31 (14), 3934–3950. <https://doi.org/10.1111/mec.16549>

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). Environmental DNA: For Biodiversity Research and Monitoring. In *Environmental DNA: For Biodiversity Research and Monitoring* . <https://doi.org/10.1093/oso/9780198767220.001.0001>

Torti, A., Lever, M. A., & Jorgensen, B. B. (2015). Origin, dynamics, and implications of extracellular DNA pools in marine sediments. *Marine Genomics* , 24 Pt 3 , 185–196. <https://doi.org/10.1016/j.margen.2015.08.007>

Trego, A., Keating, C., Nzeteu, C., Graham, A., O’Flaherty, V., & Ijaz, U. Z. (2022). Beyond Basic Diversity Estimates—Analytical Tools for Mechanistic Interpretations of Amplicon Sequencing Data. *Microorganisms* , 10 (10), Article 10. <https://doi.org/10.3390/microorganisms10101961>

Vos, M., Wolf, A. B., Jennings, S. J., & Kowalchuk, G. A. (2013). Micro-scale determinants of bacterial diversity in soil. *FEMS Microbiology Reviews* , 37 (6), 936–954. <https://doi.org/10.1111/1574-6976.12023>

Warton, D. I., Blanchet, F. G., O’Hara, R. B., Ovaskainen, O., Taskinen, S., Walker, S. C., & Hui, F. K. C. (2015). So many variables: Joint modeling in community ecology. *Trends in Ecology & Evolution* , 30 (12), 766–779. <https://doi.org/10.1016/j.tree.2015.09.007>

Warton, D. I., Wright, S. T., & Wang, Y. (2012). Distance-based multivariate analyses confound location and dispersion effects. *Methods in Ecology and Evolution* , 3 (1), 89–101. <https://doi.org/10.1111/j.2041-210X.2011.00127.x>

West, A. G., Digby, A., Santure, A. W., Guhlin, J. G., Dearden, P., Kākāpō Recovery Team, Taylor, M. W., & Urban, L. (2023). Capturing species-wide diversity of the gut microbiota and its relationship with genomic variation in the critically endangered kākāpō. *Molecular Ecology* , 32 (15), 4224–4241. <https://doi.org/10.1111/mec.16999>

Wille, M., Eden, J.-S., Shi, M., Klaassen, M., Hurt, A. C., & Holmes, E. C. (2018). Virus–virus interactions and host ecology are associated with RNA virome structure in wild birds. *Molecular Ecology* , 27 (24), 5263–5278. <https://doi.org/10.1111/mec.14918>

Willoughby, J. R., Wijayawardena, B. K., Sundaram, M., Swihart, R. K., & DeWoody, J. A. (2016). The importance of including imperfect detection models in eDNA experimental design. *Molecular Ecology Resources* , 16 (4), 837–844. <https://doi.org/10.1111/1755-0998.12531>

Wood, G., Steinberg, P. D., Campbell, A. H., Vergés, A., Coleman, M. A., & Marzinelli, E. M. (2022). Host genetics, phenotype and geography structure the microbiome of a foundational seaweed. *Molecular Ecology*

, 31 (7), 2189–2206. <https://doi.org/10.1111/mec.16378>

Woodcock, S., van der Gast, C. J., Bell, T., Lunn, M., Curtis, T. P., Head, I. M., & Sloan, W. T. (2007). Neutral assembly of bacterial communities. *FEMS Microbiology Ecology*, 62 (2), 171–180. <https://doi.org/10.1111/j.1574-6941.2007.00379.x>

Wu, L., Yang, F., Feng, J., Tao, X., Qi, Q., Wang, C., Schuur, E. A. G., Bracho, R., Huang, Y., Cole, J. R., Tiedje, J. M., & Zhou, J. (2022). Permafrost thaw with warming reduces microbial metabolic capacities in subsurface soils. *Molecular Ecology*, 31 (5), 1403–1415. <https://doi.org/10.1111/mec.16319>

Zapala, M. A., & Schork, N. J. (2012). Statistical properties of multivariate distance matrix regression for high-dimensional data analysis. *Frontiers in Genetics*, 3, 190. <https://doi.org/10.3389/fgene.2012.00190>

Zhang, Z., Geng, J., Tang, X., Fan, H., Xu, J., Wen, X., Ma, Z. (Sam), & Shi, P. (2014). Spatial heterogeneity and co-occurrence patterns of human mucosal-associated intestinal microbiota. *The ISME Journal*, 8 (4), Article 4. <https://doi.org/10.1038/ismej.2013.185>

Zinger, L., Bonin, A., Alsos, I. G., Bálint, M., Bik, H., Boyer, F., Chariton, A. A., Creer, S., Coissac, E., Deagle, B. E., De Barba, M., Dickie, I. A., Dumbrell, A. J., Ficetola, G. F., Fierer, N., Fumagalli, L., Gilbert, M. T. P., Jarman, S., Jumpponen, A., ... Taberlet, P. (2019). DNA metabarcoding—Need for robust experimental designs to draw sound ecological conclusions. *Molecular Ecology*, 28 (8), 1857–1862. <https://doi.org/10.1111/mec.15060>