

Human Monkeypox: A Potential Public Health Hazard

Amit Prasad Rath¹ and Chaitnya Prasad Meher¹

¹Centurion University of Technology and Management School of Management

April 26, 2024

Abstract

Human monkeypox (MPX) is a zoonose variola virus, causing widespread infection in humans since the 1980 eradication of smallpox. Unvaccinated individuals are vulnerable to MPXV infection. The virus's increased effectiveness in human-to-human spread, biological terrorism warnings, and increased scavenger populations make it a potential public health hazard due to its widespread prevalence and potential for scavenger populations.

Human Monkeypox: A Potential Public Health Hazard

Amit Prasad Rath^{1*}, Chaitnya Prasad Meher¹

Centurion University, Odisha, India

*Corresponding Author- amitrath222@gmail.com

Abstract :

Human monkeypox (MPX) is a zoonose variola virus, causing widespread infection in humans since the 1980 eradication of smallpox. Unvaccinated individuals are vulnerable to MPXV infection. The virus's increased effectiveness in human-to-human spread, biological terrorism warnings, and increased scavenger populations make it a potential public health hazard due to its widespread prevalence and potential for scavenger populations.

Key words : Pox virus, virology, covid, infection, isolation, disease

Introduction:

Human mpx is a zoonose variola virus like sickness affected by the causative agent pathogen, a member of the Orthopoxvirus genus. Variola virus is as well included in this genus (the causative agent of smallpox) Since the in 1980 eradication of smallpox, monkeypox has progressively arrive as the most important opxv(orthopox virus) in terms of community health, owing to its prevalence in humans Unvaccinated people are vulnerable to MPXV infection. Furthermore, the enhanced effectiveness of MPXV human-to-human Spread, the biological terrorism warning of smpx and MpXv, and the rise in the noticed numeral of scavenger(animal) masses all point to mPxv as a possible huge-range civic well-being hazard Human monkeypox was found in 1970, after the discovery of occasional cases of a comparable disease in rural parts of the Democratic Republic of the Congo [DRC]. Since then, the epidemiology and clinical aspects of the condition have been intensively studied, but no effective countermeasures have been implemented Human instances of monkeypox were caused by dealings with infectious rodents that had been stored or convey alongside African cavy(rodents) bring in from gold coast (Ghana) Monkey pox is meticulously correlated to smpx, and the smallpox preparation is assume to be defensive in contradiction of Mpx . Mpxv was initial revealed in baboons. Since Monkey-pox has equally zoonotic and homosapien to homosapien spread, the first case was testified in1970 DRc(Democratic Republic of the Congo) , and it has expand to several realms both within and apart from of Africa.Zoonosis

spread occurs through straight interaction through infected animals' plasma, physique liquids . Ineffectively roasted substance may also be a factor. Human to anthropoid transference occurs through undeviating by way of bodily fluids infected people's epidermis sores, or coming into exposure with tainted particles by a contaminate individual human beings. The West and Central Africa are indigenous for monkeypox microbe however following its first detection in the DrC- congo in 1970, the number of sufferer has increased to the point where it is now present in several African and European regions. According to research, the rise in Monkeypox case scenarios the human species perhaps assign to the cessation of smallpox immunize, which generally provides cross shield opposed to Monkeypox thus the term "monkeypox." The word is misleading cuz the virus's big animal zone have been discovered in cavy(rodents) such as chipmunk and large gopher, both of which are chase for nourishment Smallpox does not cause lymphadenopathy, whereas monkeypox does. The affliction may last for 28 days before sarcomas on the epidermis go away. MpXv was discovered In 1958 throughout a virulent disease of monkeypox with inside the Asian monkey "Macaca fascicularis" which became used for polio vaccine studies at a Copenhagen, Denmark beast provision in August 1970 the precedent of hmpxv was discovered at Basankusu sanatorium in Zaire's Ecuador region, when a nine-month-vintage boy evolved a smallpox-like fragility. The Collaborating Center of the WHO at the investigate academy for Viral procure in Moskva confirmed that monkeypox virus was the cause of the boy's illness. Even though the arena is existing dealing with the COVID-19 ailment, the emergence of a brand new outburst precipitated by the m.pox virus has citizen health ceremonial is concerned about whether may it will pose a new warning monkeypox virus is a double stranded Deoxyribonucleic acid virus that belongs to orthopox viruses class, which also includes variola, CPX(cowpox) afterward the abolition of small-pox monkey-pox has emerged as the most common opxv influence human section Monkeypox is now indigenous in Western and Central African countries which have experienced a rising of monkeypox explosion in the last 10yr.Since 2016, there have been more validate cases of monkeypox than in the previous 40 years.

History :-

MPX is due to the zoonotic monkey pox microbe a part of the Orthopoxvirus aptitude the 1st anthropoid Mpxv incident turned into stated in 1970 withinside DRc(Democratic Republic of Congo), which remains Africa's place with the best degree of endemicity

MPX research has largely gone unnoticed. Epidermis lesions, pyrexia(fever), and enlarged lymph nodes are common clinical manifestations. MPX is commonly self-limiting however excessive instances can happen, and a incident death rate of 1 to 10% has been mentioned from Africa with infections from the Central African viral clade being associated with a higher case fatality rate

Orthopoxvirus is a member of the Poxviridae subfamily Chordopoxvirinae. Family is a member of the order Chitovirales, which is a subclass of the class Pokkesviricetes. This class is a member of the phylum Nucleocytoviricota, the dominion Bamfordvirae

Members of the subdivision OpXv and also additional extremity of the family Pox viridae that affect rabbits, insects, birds and goats cause disease in humans and animals.

Drug Absorption, Distribution, Metabolism, Excretion Analysis :

Simplified Molecular Input Line Entry System is a notation system used to represent the structure of molecules as a linear string of characters. It provides a compact and standardized way to describe molecular structures, making it easier to input and process them in computational tools. The ligands (phytochemicals) were initially screened to ensure they adhere to Lipinski's rule of five. This rule is a guideline used in drug discovery to assess the likelihood of a compound's oral bioavailability. It evaluates properties such as molecular weight, lipophilicity, hydrogen bonding, and rotatable bonds. After the screening, the molecular and physicochemical properties of the phytochemicals were investigated using the Swiss ADME tool. Swiss ADME is a web based platform that provides various computational models and algorithms to predict molecular properties, including solubility, lipophilicity, and drug-likeness. To further analyse the phytochemicals, they were converted from the Protein Data Bank to the standard SMILES format. This conversion allows for easier input and compatibility with various computational tools. The SMILES-formatted phytochemi-

cals were then uploaded to the Swiss ADME prediction tool to predict their drug absorption, distribution, metabolism, excretion, and toxicity profiles. This information is essential in drug discovery and development, as it helps assess the potential efficacy and safety of the compounds. Additionally, the algorithm based pharmacokinetics , pkCSM was utilized to computationally predict the physicochemical and pharmacological properties of the phytochemicals. This server employs various algorithms and models to estimate parameters related to the absorption, distribution, metabolism, excretion, and toxicity of small molecules. Overall, the use of SMILES format, along with computational tools enables researchers to efficiently analyse and predict the properties of phytochemicals, aiding in the drug discovery and development process.

Symptom

- (1) Lethargy
- (2) Asthenia
- (3) Lymph node swelling
- (4) Seizures
- (5) Confusion
- (6) Joint pain and/or swelling
- (7) Diarrhea
- (8) Wheezing
- (9) Ear pain
- (10) Conjunctivitis
- (11) Abdominal pain
- (12) Chest pain
- (13) Dysphagia
- (14) Stiff neck
- (15) Shortness of breath
- (16) Mouth sores
- (17) Back pain
- (18) Nasal congestion
- (19) Nausea and/or vomiting
- (20) Cough
- (21) Sweats
- (22) Myalgia
- (23) Sore throat
- (24) Headache
- (25) Adenopathy
- (26) Chills and/or rigors
- (27) Fever
- (28) Rash

Diagnosis:

Viral DNA analysis using technique of polymerase chain reaction (PCR) is the preferred laboratory diagnostic for mpox. By firmly wiping skin, liquid or the outer layer from the rash, the best clinical examination may be taken. If there are no cutaneous lesions testing can be done on oropharyngeal, anal or rectal swabs. Not recommended is examinations of vital fluid. The inability to differentiate between different orthopox viruses may render them ineffective.

The allergic reaction initially appears as unpleasant, flat, red pimples. These lumps develop into blisters that ooze pus. Blisters that develop eventually harden and peel off. may take 2 to 4 weeks

(i) RT - PCR

The best screening specimens include dermatological lesions, pustule and blister fluid, and dry outermost layer etc

(ii) WGS

Wgs is highly expensive technology so this technology is not use

(iii) serology

This technology is not recommended due to the cross-reactive with other pox virus

Clinical examination :-

skin, vaginal mucosa, and oral mucosa

Symptomatic treatment :-

-Skin irritation and genital lesions

- Fever > 38°C

Real time PCR :-

- Any suspicious skin or mucosal lesion should be swabbed.

Monkey pox negative :-

- Stop monitoring

Monkey pox positive :-

Hospitalization in a tertiary or recognised facility

WHO guideline :-

1. Severe - 100 to 250 skin lesions
2. Moderate - 25 to 99 skin lesions
3. Mild - less than 25 skin lesions

Fetal :-

Depending on gestational age, fetal evaluation corticosteroid use for foetal lung maturation

Delivery :-

The virus can also spread during pregnancy to the foetus, during or after birth via skin-to-skin contact, or from a parent infected with monkeypox to an infant or child via intimate contact.

Newborn :-

- The newborn should be cleaned as soon as possible

- Real-time PCR for monkey pox in newborn baby

According to lab studies:-

Electron microscopy of monkeypox rash specimens displays distinctive big slab(brick)-formed elements. In vesicular liquids, the mulberry arrangement predominates, with mulberry bulge visible and a lack of PTA (phosphor tungstic acid) stain perforation into the constituent part

Prevalence research :-

In 1981, surveys of people without vaccination marks were done in Congo, Zaire, Côte d'Ivoire, Sierra Leone (regions of Central and West Africa affected by monkeypox detected in 1970s). By HAI (hemagglutination-inhibition Testing), an orthopoxvirus general Elisa test, 663/3460 (19.2%) of DRC (then Zaire) samples were seropositive. 178 of the hemagglutination forbid testing-positive sera were verified using the Radio immuno assay (RIA) test, which is more specific to monkeypox. 27 (15%) of the 178 sera verified were +ve for "monkeypox."

Disease prognosticate :

Modeling study results based on human epidemiology, as well as limited ecological educations, were used to determine monkey pox could accurately spread among people and exchange spx(smallpox) as a sever, entomb social communicable pathogen. These scrutiny also influenced the verdict not to reintroduce "smallpox" inoculation. By means of records from the 1981 to 1986 research.

Difficulty:-

There were secondary bacterial infections recorded. Eyelid ulceration was observed in 4 percent of not vaccinated case patients and 1% of initially vaccinate case forbearing(patients). Bronchial pneumonia & respiratory illness were detected late in the sickness (were thought to be caused by bacteria); 19 of 34 unvaccinated patients with this difficulty pass away. Broncho-pneumonia collide with 34/270 unvaccinated patients and 2/43 earlier s.pox-immunized stoic(patient). Additional problems combined nausea and diarrhoea, which resulted in water loss.

Oropharynx:-

Oropharynx abnormalities were originate in 76% of unvaccinated and 47% of initially variola-immunized monkeypox patients.

upper respiratory :-

palatine tonsils and oral ulcers have been noticed. tussis (cough) was reported in 39% of non vaccinated patients and 14% of recently immunised patients.

ocular problem:-

Keratitis and inflammation of the eyelids. were observed in 30% of unimmunized patients and 7% of earlier immunised patients.

Prevention :

Due to the shortcomings of the currently available smallpox vaccines, research into alternative therapeutics like as immunoglobulin and antiviral medicines is critical in stoping excessive or deadly oPxV infection in immunodeficient individuals.

Who don't need therapy intended for medicinal reasons can be solitude in the home and different non healthcare surroundings the usage of suitable infect control measures. Prevention of infection transmission via inhaling and contact path is also essential in the home.

(1) Type of household situation, such as at home, an apartment complex, a direct provision centre, a hostel, etc.

(2) Unable to keep away from contact with immunocompromised individuals, child-bearing, and children lower than the age of 13 years.

Hand Hygiene :

washing with cleaning soapsuds and H₂O or the use of an spirits-primarily based hand polish execute by sick individuals and family associates after touching clothing linens, lesion material and ecological surfaces that could have come into touch with mutilation (lesion) substances

Lavatory:

Cleaning home settings need to be done in the order listed below:-

- Before entering the room, contaminated clothing and linens should be collected.
- cleaned. Personal apparel or line have to now no longer be shaken or treated in any way that could dispel contagious particles
- element that have come into direct touch with an infected person's skin and are difficult to wash in a house wash appliance, such as pillows, comforters and coverlets, can be positioned in a sealed and bag.
- Handling dirty laundry, which include towels, private clothing, and bedding should be done with caution to keep away from direct touch with infected material. One-use gauntlet(gloves) have to be shabby and hands must be thoroughly wiped clean later eliminating gloves the. Wherever possible, this have to be done with the aid of using the showed case.
- To avoid the spread of virus particles and skin scales, home linen in a not reusable carrier bag before transferring it to the washing machine.
- Clothing and linens that have been contaminated must be laundered at 60°C on an prolonged rotation. Do not excess the washing appliance (goal for hemi or three quarters full) and dodge smaller 'economy series' (individuals that use less H₂O and save vitality) until the distinct has improved completely.
- Confirmed MPX cases must do their private dirty washing whenever feasible, keeping their washing matters apart after the rest of the house's laundry and washing them with their regular cleaner, per producer's directions.
- Items that have been washed should not be located in zones where they can become re-infect during the scrubbing method.
- If a person does not have a washing device, they can hand rinse their clothes with warm liquid and regular cleansing agent. This may work better in a big hand basin or plunge bath when finished, it is critical to hygienic and sanitize all outsides with disposable gloves. When cleaning these surfaces with bleach, use extra caution.

Environmental Cleaning

- Steam cleaning can be used on carpets, curtains, and other soft furniture.
- Individuals should clean and dry their specific used saucers and intake appliance, then if they have one, use a dish washer with warm H₂O (above 60oC) and detergent. If this is not possible, then eating paraphernalia(utensil) should be mop with regular dishwashing liquid and hot water and air dried.
- If a person takes lacerations on their fingers and does not have access to a wash, they must be instructed to wash up with solitary-use only not reusable gloves. Recyclable gloves should not be shared and have to be thrown out once the segregation period has ended.
- Carpets, for example, can be washed with a High Efficiency Particle Arresting filtered vacuum cleanser (if one is accessible); however, carefulness Must be obtained when arranging of the dust buster contents to

diminish dust particle dispersal. Vacuum cleaner unwanted should be emptied wisely into a not reusable trash basket.

- As an added safety measure, all not reusable trash bags have to be located in a additional throwaway carrier and firmly sealed before being willing of as normal with household discarded. All garbage container should be strongly put in storage up to they are collected. Unused should not be placed in reprocessing containers until the self-isolation period has completed.

sexual activity

Early results from research indicate that being a young guy, having intercourse with other males, partaking in dangerous behaviours and activities, such as condom less sex, and sero positive are risk factors for monkey pox. HIV testing positive and a history of STIs, including syphilis

Given the uncertainty surrounding monkeypox virus sexual transmission via sperm or different sexual frame fluid, showed instances who want to renew sexual action after self-solitude has over are suggested to apply a protection for 84days afterwards the impetuous has scraped done and layers have dropped off. It is a preventive measure toward decrease the possibility of infection distribution to others. This is a cautionary measure. This advice may change as new proof becomes available.

Personal Protective Equipment :

(1) All confirmed cases, especially those with respiratory symptoms, should wear a medical grade (surgical) mask

(2) The somebody kind intended for or helping the individual with MPX have a duty to wear one-use gloves

Safety for pregnant woman :-

For the reason that of the post-COVID-19 edge regenerating and tourism between nations currently undergoing an epidemic, the risk of contagion in pregnant females is high. Human infections with monkeypox and smallpox (both orthopoxviruses) transmit a high danger of highly inherited contamination, maternal illness, miscarriage, and death

fertilization woman with monkeypox :-

- Traveled to an impacted nation within the past 21 days

- Sexual intercourse with a proven case of monkeypox

Pharmacotherapy :

Monkeypox does not have a specific treatment. The main recommendations remain supportive care, symptomatic management, and mitigation of subordinate bacterial contamination Prior immunisation by the help of smallpox serum appears to have a defensive impact in opposition to monkey pox virus and can enhance scientific manifestations of infection. The United State SNS (Strategic National Stockpile) presently carries 3 smallpox vaccines: Jynneos (additionally called as, Imvanex, Imvamune, Mva-Bn) and Acam2000 are licenced for smpx, and the APSV (Aventis-pasteur-smallpox-Vaccine) could be used for smallpox under an IND (investigational-new-drug) procedure

JYNNEOSTM is a alive biological inoculation made from a improved strain of vabn (vaccinia-Ankara-Bavarian-Nordic) that was approved by the UnitedState FDA (Food and Drug Administration) in September 2019.

ACAM2000 consists of stay vaccinia germ also. It become authorized in August 2007 in fda but was removed by the producer because it is intended for effective immunisation antagonist of smallpox affliction in people who have been determined to be at high risk of infection.

APSV (Aventis-Pasteur-SmallpoxVaccine) is a duplication-capable vaccine that can be applied with an Ind or contingency Usage Approval. However, it is unknown whether this serum could be apply to prevent mPxV.

Vaccine :

Vulnerability vaccination of previously unvaccinated individuals against smallpox: deliver a course of two doses, with at least a 28-day interval between doses, for instance :

- (1) 0.5ml MVA-BN vaccine initial dose, then
- (2) A second 0.5ml dose of MVA-BN vaccine should be administered at least 28 days after the first dose.

Individuals of any age should be given a single 0.5ml dose of MVA-BN vaccine after being exposed.

The vaccine should be administered via deep subcutaneous injection

Drugs use in Monkey pox virus:

Treatments

- (1) Tecovirimat :

Route : IV

Dose : Adults: 600 mg TID for 14 days

Pediatrics : 13kg to 25kg - 200 mg BID for 14 days

Side effects : Infusion-site reactions , vomiting

Tecovirimat showed promising antiviral activity against both Monkeypox virus and a virus similar to smallpox in rabbits in subsequent investigations, the pharmacokinetics (how the drug is absorbed, distributed, metabolized, and excreted) and safety of tecovirimat were assessed in a clinical trial involving 361 healthy individuals. These participants were randomly assigned to receive an oral dose of 600 mg of tecovirimat twice a day. The results of the study indicated that tecovirimat was four times more effective in non-human primates compared to humans. However, no significant safety concerns or warning signs of danger were observed during the trial. The adverse effects of tecovirimat were minimal. These findings suggest that tecovirimat has potential as an effective antiviral agent against orthopoxviruses, including the Vaccinia virus, cowpox virus, Monkeypox virus, and a virus similar to smallpox. Further research and clinical trials are needed to fully understand its efficacy and safety in humans.

- (2) Brincidofovir :


Route : Tablet , syrup

Dose : Adult - >48 kg – 200mg weekly for two days

Pediatrics : >10kg – 4mg/kg – syrup

Side effects : Diarrhea, adomen cramp

Chemical structure:



(Brincidofovir)

Controlling:

Itchy skin –

(i) Clean with Antiseptic

(ii) If secondary infection may develop into the skin then Antibacterial drug may be prescribed

Genitalia ulcer –

(i) hip bath

Mouth ulcer –

(i) Anti-inflammatory drugs may be used

Eye inflammation –

(i) Chloramphenicol eye drop may be used

(ii) highly effective Antibiotic may be used

(iii) Antipyretic drug are used in this condition

Conclusion:

Most cases of monkeypox are found in central and western Africa. The ailment is a typical zoonosis, unlike smallpox, in that most cases come from having be in touch with with an infected animal.

REFERENCE

(1) Adesola Yinka-Ogunleye, Olusola Aruna, Mahmood Dalhat, Dimie Ogoina, Andrea McCollum, Yahyah Disu, Ibrahim Mamadu, Afolabi Akinpelu,

Adama Ahmad, Joel Burga, Adolphe Ndoreraho, Edouard Nkuzimana, Lamin Manneh, Amina Mohammed, Olawunmi Adeoye, Daniel Tom-Aba,

Bernard Silenou, Oladipupo Ipadeola, Muhammad Saleh, Ayodele Adeyemo, Ifeoma Nwadiutor, Neni Aworabhi, Patience Uke, Doris John,

Paul Wakama, Mary Reynolds, Matthew R Mauldin, Jeffrey Doty, Kimberly Wilkins, Joy Musa, Asheena Khalakdina, Adebayo Adedeji, Nwando Mba, Olubunmi Ojo, Gerard Krause*, Chikwe Ihekweazu* Published Online July 5, 2019 , DOI:([http://dx.doi.org/10.1016/S1473-3099\(19\)30281-6](http://dx.doi.org/10.1016/S1473-3099(19)30281-6)), Outbreak of human monkeypox in Nigeria in 2017–18:a clinical and epidemiological report, 1/8

(2) Julie Dyall, Reed F. Johnson, Dar-Yeong Chen, Louis Huzella, Dan R. Ragland, Daniel J. Mollura, Russell Byrum, Richard C. Reba, Gerald Jennings, Peter B. Jahrling, Joseph E. Blaney, and Jason Paragas , Published by Oxford University , DOI: 10.1093/infdis/jir663 , Molecular Imaging of Monkeypox , 1/10

(3) Brett W. Petersen, Kabamba, J., McCollum, A.M., Lushima, R.S., Wemakoy, E.O., Muyembe Tamfum, J.J., Nguete, B., Reynolds, M.G., Vaccinating against monkeypox in the Democratic Republic of the Congo, Antiviral Research (2018), DOI:<https://doi.org/10.1016/j.antiviral.2018.11.004> , 1/23

(4) David A Kulesh, Bonnie M Loveless, David Norwood, Jeffrey Garrison, Chris A Whitehouse, Chris Hartmann, Eric Mucker, David Miller, Leonard P Wasieloski Jr, John Huggins, Gregory Huhn4, Lori L Miser, Carroll Imig, Mark Martinez, Tom Larsen, Cynthia A Rossi and George V Ludwig , published online 21 June 2004 , Monkeypox virus detection in rodents using real-time 30-minor groove binder TaqMans assays on the Roche LightCycler , 1/9

(5) Mansoor Ahmed, Haseena Naseer, Mateen Arshad, Afnan Ahmad , Monkeypox in 2022: A new threat in developing , Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd 7 June 2022 , DOI:<https://doi.org/10.1016/j.amsu.2022.103975> , 1/3

(6) Ibrahim Idris a,* , Ridwan Olamilekan Adesola , Monkeypox virus

United Kingdom Endemic and surveillance , Publishing services by Elsevier B.V , DOI:<https://doi.org/10.1016/j.bj.2022.07.007> , 1/3

(7) Eskild Petersen, MD, DMSc, DTMH , Anu Kantele, MD, PhD, Marion Koopmans, DVM, PhD Danny Asogun, MBBS, FWACP, Adesola Yinka Ogunleye, BDS, MPH, Chikwe Ihekweazu, MBBS, MPH, FFPH, Alimuddin Zumla, FRCP, PhD, FRCPATH , Human Monkeypox Epidemiologic and Clinical Characteristics, Diagnosis, and Prevention , Published by Elsevier Ltd , DOI:<https://doi.org/10.1016/j.idc.2019.03.001> , 1/17

(8) Harapan Harapan, Abdul M. Setiawan, Amanda Yufika, Samsul Anwar, Sri

Wahyuni, Febrivan W. Asrizal, Muhammad R. Sufri, Reza P. Putra, Nanda P.

Wijayanti, Salwiyadi Salwiyadi, Razi Maulana, Afriyani Khusna, Ina Nusrina,

Muhammad Shidiq, Devi Fitriani, Muharrir Muharrir, Cut A. Husna, Fitria

Yusri, Reza Maulana, Mohd Andalas, Abram L. Wagner & Mudatsir Mudatsir , Knowledge of human monkeypox viral infection among general practitioners: across-sectional study in Indonesia , DOI:<https://doi.org/10.1080/20477724.2020.1743037> , 1/9

(9) Z. Jezek, M. Szczeniowski, K. M. Paluku, and M. Mutombo , Human Monkeypox: Clinical Features of 282 Patients , <http://jid.oxfordjournals.org/> at Russian Archive on January 8/2014 , 1/6

(10) John G. Rizk · Giuseppe Lippi · Brandon M. Henry · Donald N. Forthal · Youssef Rizk , Prevention and Treatment of Monkeypox , <https://doi.org/10.1007/s40265-022-01742-y> , 1/7

(11) Ziyue Wang, Pinkus Tober-Lau , Vadim Farztdinov, Oliver Lemke, Torsten Schwecke, Sarah Steinbrecher, Julia Muenzner, Helene Kriedemann , Leif Erik Sander , Johannes Hartl , Michael M(eurou)lleder, Markus Ralser & Florian Kurth, The human host response to monkeypox infection: a proteomic case series study , Published under the terms of the CC BY 4.0 license , DOI 10.15252/emmm.202216643 , 1/11

- (12) Varea Costello, Madeleine Sowash, Aahana Gaur, Michael Cardis, Helena Pasieka, Glenn Wortmann, Sheena Ramdeen ,Emerging Infectious Diseases , Vol. 28, No. 5 , DOI:<https://doi.org/10.3201/eid2805.220292> , 1/4
- (13) Vicente A. Benites-Zapata, Juan R. Ulloque-Badaracco, Esteban A. Alarcon-Braga, Enrique A. Hernandez-Bustamante, Melany D.Mosquera-Rojas, D. Katterine Bonilla-Aldana^{5,6} and Alfonso J.Rodriguez-Morales , Annals of Clinical Microbiology and Antimicrobials , DOI:- <https://doi.org/10.1186/s12941-022-00527-1> , 1/18.
- (14) Gregory D. Huhn, Audrey M. Bauer, Krista Yorita, Mary Beth Graham, James Sejvar, Anna Likos,² Inger K. Damon, Mary G. Reynolds, and Matthew J. Kuehnert , Clinical Characteristics of Human Monkeypox, and Risk Factors for Severe Disease , <https://academic.oup.com/cid/article/41/12/1742/344953> , 1/10
- (15) Professor Sir Gregor Smith , Professor Alex McMahon, Professor Alison Strath , MONKEYPOX (MPX) PRE AND POST EXPOSURE VACCINATION , Pulished - 18 July 2022
- (16) Katy Brown and Peter A. Leggat , Human Monkeypox: Current State of Knowledge and Implications for the Future , Published: 20 December 2016 , DOI:[10.3390/tropicalmed1010008](https://doi.org/10.1093/tropicalmed/taab008) , 1/13
- (17) Interim Public Health Guidance for the Management of Monkeypox Cases and their Contacts , Version 1.8 , 1/21
- (18) Guidelines for pregnant individuals with monkeypox virus exposure , Published Online June 21, 2022 [https://doi.org/10.1016/S0140-6736\(22\)01063-7](https://doi.org/10.1016/S0140-6736(22)01063-7) , 1/ 2
- (19) Inger K. Damon , Status of human monkeypox: clinical disease, epidemiology and research , Published by Elsevier Ltd , <http://www.elsevier.com/locate/vaccine> , 1/6
- (20) AbdulRahman A. Saied, Manish Dhawan , Asmaa . Metwally , Mathumalar Loganathan Fahrni , Priyanka Choudhary and Om Prakash Choudhary , Vaccines 2022, 10, 2091. <https://doi.org/10.3390/vaccines10122091> .
- (21) Pallavi Gulati , Jatin Chadha , Kusum Harjai and Sandeepa Singh ,Targeting envelope proteins of poxviruses to repurpose phytochemicals against monkeypox: An in silico investigation , <https://doi.org/10.3389/fmicb.2022.1073419>
- (22) Adam Sherwat, M.D., John T. Brooks, M.D., Debra Birnkrant, M.D., and Peter Kim, M.D , Tecovirimat and the Treatment of Monkeypox — Past, Present, and Future Considerations , DOI: 10.1056/NEJM2210125
- (23) GUIDELINES FOR MANAGEMENT OF MONKEYPOX DISEASE Ministry of Health and Family Welfare GOVERNMENT OF INDIA