

Table 1 General and clinical characteristics of the study population.

Abbreviation: ED: emergency department. CT: computerized tomography; MRI: Magnetic Resonance Imaging; ICU: Intensive Care Unit.

Note: Variables are expressed as frequencies (percentages), arithmetic means (\pm SD) or median (interquartile range, IQR).

	No. of children (30)
Age, median (IQR), years-months	8yy 7mm (5yy 6mm -13yy 9mm)
Males, No. (%)	18 (60)
Transfer from another ED, No. (%)	5 (16.7)
Symptoms and signs	
Chest pain, No. (%)	1 (3.3)
Cough, No. (%)	12 (40)
Fever, No. (%)	19 (63.3)
Fever, median(IQR), number days before ED visit	2 (0-4)
Exertional dyspnea, No. (%)	4 (13.3)
Thoracic pain, No. (%)	1 (3.3)
Respiratory distress, No. (%)	0 (0)
Seizure, No. (%)	1 (3.3)

Symptoms of upper respiratory tract infection, No. (%)	30 (100)
Arthralgia, No. (%)	5 (16.7)
Headache, No. (%)	4 (13.3)
Diarrhea, No. (%)	3 (10)
Vomiting, No. (%)	0 (0)
Imaging and other	
Oxygen saturation in ED, mean (SD), %	98.8 (1.05)
Period of hospitalization, median (IQR), day	10 (7-12)
Chest x-ray, No. (%)	12 (40)
Chest CT, No. (%)	4 (13.3)
Chest MRI, No. (%)	2 (6.7)
Oxygen therapy, No. (%)	0 (0)
Cases in ICU, No. (%)	1 (3.3)
Therapy	
Nothing, No. (%)	14 (46.7)
Azithromycin, No. (%)	2 (6.7)
Hydroxychloroquine and Azithromycin, No. (%)	7 (23.3)
Other therapy, No. (%)	7 (23.3)