

1    **Temperature dependence of clumped isotopes ( $\Delta_{47}$ ) in aragonite**

2    Niels J. de Winter<sup>1,2</sup>, Rob Witbaard<sup>3</sup>, Ilja J. Kocken<sup>2</sup>, Inigo A. Müller<sup>4</sup>, Jingjing Guo<sup>2</sup>, Barbara  
3    Goudsmit<sup>2,3</sup>, Martin Ziegler<sup>2</sup>

4

5    <sup>1</sup>Analytical, Environmental and Geochemistry Group, Vrije Universiteit Brussel, Belgium

6    <sup>2</sup>Dept. of Earth Sciences, Utrecht University, the Netherlands

7    <sup>3</sup>Dept. of Estuarine and Delta Systems, Royal Netherlands Institute for Sea Research, the  
8    Netherlands.

9    <sup>4</sup>Department of Earth Science, University of Geneva, Switzerland

10

## Key points

- Precise control on carbonate formation temperatures enables more accurate clumped isotope-temperature calibrations
- Isotopic ordering and acid fractionation in aragonite have a similar temperature dependence as in calcite, enabling combined calibrations
- The  $\Delta_{47}-\frac{1}{T^2}$  relation in carbonates is non-linear, including hot calibration data offsets the calibration in the cold temperature range

## Abstract

Clumped isotope thermometry can independently constrain the formation temperatures of carbonates, but a lack of precisely temperature-controlled calibration samples limits its application on aragonites. To address this issue, we present clumped isotope compositions of aragonitic bivalve shells grown under highly controlled temperatures (1–18°C), which we combine with clumped isotope data from natural and synthetic aragonites from a wide range of temperatures (1–850°C). We observe no discernible offset in clumped isotope values between aragonitic foraminifera, mollusks, and abiogenic aragonites or between aragonites and calcites, eliminating the need for a mineral-specific calibration or acid fractionation factor. However, due to non-linear behavior of the clumped isotope thermometer, including high-temperature (>100°C) datapoints in linear clumped isotope calibrations causes them to underestimate temperatures of cold (1–18°C) carbonates by  $2.7 \pm 2.0^\circ\text{C}$  (95% confidence level). Therefore, clumped isotope-based paleoclimate reconstructions should be calibrated using samples with well constrained formation temperatures close to those of the samples.

## Plain language summary

Clumped isotope analysis is a highly accurate method for reconstructing temperatures in Earth's past climate from calcium carbonate fossils of calcifying organisms. Unfortunately, calibration studies so far were predominantly based on samples of calcite, a common calcium carbonate mineral. It is therefore unknown whether these clumped isotope calibrations yield accurate temperature reconstructions when applied to aragonite, another carbonate mineral which corals and many shells consist of. Therefore, we grew mollusks that build their shell out of aragonite in a lab at constant water temperatures to test the clumped isotope method on aragonitic shells. We find no significant difference in the temperature sensitivity of the method between our aragonites and the previous calibrations and show that the temperature calibration can be improved by combining data from different minerals. However, we find subtle differences in the temperature dependence of clumped isotopes between hot ( $>100^{\circ}\text{C}$ ) carbonates and cold ( $<30^{\circ}\text{C}$ ) carbonates, which cause previous calibrations to underestimate temperatures of colder carbonates. We conclude that using carbonate samples grown at temperatures close to the temperatures of the samples used in climate reconstructions can eliminate a bias of  $2.7^{\circ}\text{C}$ , resulting in more accurate reconstructions of past temperatures.

## **Keywords**

Clumped isotopes, aragonite, paleoclimate, mollusk, temperature

## **1. Introduction**

Since its first applications (e.g. Schauble et al., 2003; Wang et al., 2004; Ghosh et al., 2006), carbonate clumped isotope analysis has developed into a valuable tool for paleothermometry in the geosciences. Clumped isotope analysis is based on the thermodynamic principle that molecules with multiple heavy isotopes (so-called “multiply-substituted isotopologues”) have lower vibrational energies than molecules containing lighter isotopes (Urey, 1947). Consequently, the increase in system entropy at higher temperatures causes a decrease in

the occurrence of multiply-substituted isotopologues, and “clumping” of heavy isotopes within the same molecule is favored in low-energy systems (Eiler, 2007). In carbonates, this principle causes heavy carbonate ions (e.g.  $^{13}\text{C}^{18}\text{O}^{16}\text{O}_2$ ; mass 63 or  $^{12}\text{C}^{18}\text{O}_2^{16}\text{O}$ ; mass 64) to become more abundant with decreasing calcification temperatures (Ghosh et al., 2006). The distribution of these isotopologues is proportional in the  $\text{CO}_2$  gas after reaction of carbonates with acid (e.g.  $^{13}\text{C}^{18}\text{O}^{16}\text{O}$ ; mass 47 and  $^{12}\text{C}^{18}\text{O}_2$ , mass 48 respectively) and is measured with reference to the distribution of isotopologues in a fully scrambled heated  $\text{CO}_2$  gas with the same isotopic composition:

$$\Delta_{47}[\text{‰}] = \left( \frac{R^{47}}{R^{47*}} - 1 \right) \quad (1)$$

In which  $R^{47}$  is the ratio of  $\text{CO}_2$  molecules with mass 47 (predominantly  $^{13}\text{C}^{18}\text{O}^{16}\text{O}$ ) relative to  $\text{CO}_2$  with the most common mass 44 ( $^{12}\text{C}^{16}\text{O}_2$ ) in the sample, and  $R^{47*}$  is the same ratio in stochastic equilibrium (Daëron et al., 2016). This  $\Delta_{47}$  value is a measure for the degree of “clumping” in the sample which depends on its calcification temperature.

The main advantage of carbonate clumped isotope analysis over previous paleothermometers is its basis on thermodynamic principles and its independence from the chemistry of the precipitation fluid (Eiler, 2007). The latter represents an improvement over the often-used oxygen isotope paleothermometer ( $\delta^{18}\text{O}$ ), which requires knowledge of the oxygen isotope composition of the precipitation fluid ( $\delta^{18}\text{O}_w$ ; e.g. Epstein et al., 1953; Kim & O’Neil, 1997). The clumped isotope method has many applications, notably to reconstruct absolute temperature variability throughout Earth’s history (Rodríguez-Sanz et al., 2017; Henkes et al., 2018; Vickers et al., 2020a; de Winter et al., 2021a; Meckler et al., 2022; Agterhuis et al., 2022).

Inter-lab standardization of carbonate  $\Delta_{47}$  measurements has reconciled former offsets between laboratories using different  $\text{CO}_2$  preparation methods and reconciled the clumped isotope temperature calibration of calcites with the results of thermodynamic *ab initio* models (Bernasconi et al., 2018; 2021; Petersen et al., 2019; Jautzy et al., 2020). A unified linear calibration was established through re-standardized  $\Delta_{47}$  values of carbonates precipitated at a

wide range of known temperatures (0.5-1100°C; Anderson et al., 2021). This eliminates concerns over the confounding effects of differences in the origin of carbonates (e.g. biogenic vs. inorganic; Henkes et al., 2013), varying mineralization rates (Daëron et al., 2019), different acid digestion temperatures and different carbonate mineralogies (e.g. dolomite vs. calcite; Müller et al., 2019) on the clumped isotope thermometer. However, it remains unclear whether biological process (i.e. “vital effects”) influence isotopic ordering in some biogenic carbonates.

The unified calibration dataset includes only one aragonitic carbonate, insufficient to test for different clumped isotope temperature dependencies between aragonites and calcites (Anderson et al., 2021). Results of *ab initio* models suggest that such a difference between the two polymorphs may exist (Schauble et al., 2006; Guo et al., 2009) and experimental studies disagree on a difference in acid fractionation factor (AFF) between calcite and aragonite (Guo et al., 2009; Müller et al., 2019; Petersen et al., 2019). These uncertainties are confounded by the fact that most carbonates used in current calibrations are precipitated under natural circumstances with indirectly estimated or else poorly controlled temperature regimes (e.g. Kele et al., 2015; Peral et al., 2018). The potential  $\Delta_{47}$  offset between aragonite and calcite might introduce an unknown bias when using the unified temperature calibration on aragonite data (e.g. Caldarescu et al., 2021); a severe limitation given the common occurrence of aragonite in biogenic calcifiers (e.g. bivalves; Kennedy et al., 1969, gastropods; Taylor and Reid, 1990, and foraminifera; Hansen, 1979) as well as inorganic natural carbonates (e.g. speleothems; Frisia et al., 2000, and travertines; Kele et al., 2015).

This study presents new clumped isotope results from precisely temperature controlled, lab-grown aragonitic *Arctica islandica* bivalve shells. The bivalve *Arctica islandica* is a highly utilized climate archive, and a promising substrate for clumped isotope-based paleothermometry (e.g. Witbaard et al., 1997; Burchardt and Simonarson, 2003; Schöne et al., 2005; Schöne and Fiebig, 2009; Butler et al., 2013). Combined with preexisting aragonite clumped isotope data (Kluge et al., 2015; Kele et al., 2015; Müller et al., 2017; Breitenbach et al., 2018; Bernasconi et al., 2018; Piasecki et al., 2019; Caldarescu et al., 2021) standardized

to the new Intercarb-Carbon Dioxide Equilibrium Scale (I-CDES) reference frame (Bernasconi et al., 2021), our dataset resolves potential vital effects on clumped isotopes in aragonitic mollusks by comparing species and specimens grown under the same controlled conditions. This study aims to offer a detailed investigation of the clumped isotope temperature dependence in aragonites.

## 2. Materials and Methods

### 2.1 Lab grown *Arctica islandica*

*Arctica islandica* bivalves were cultured inside the lab of the Royal Netherlands Institute for Sea Research (NIOZ, Texel, the Netherlands). Specimens used for this study were grown under four different, constant, and monitored temperature regimes:  $1.1 \pm 0.2^{\circ}\text{C}$ ,  $3.2 \pm 0.3^{\circ}\text{C}$ ,  $15 \pm 0.4^{\circ}\text{C}$  and  $18 \pm 0.3^{\circ}\text{C}$  (see **Table 1**; **S9**). Details on the culturing setup are provided in **S1** and Witbaard et al. (1998). Aragonite from cleaned and dried *Arctica islandica* shells was sampled using a hand-held Dremel 3000 rotary drill at low speed equipped with a tungsten-carbide drill bit (see **S1**). Gathering enough aragonite for reliable  $\Delta_{47}$  analyses for each temperature treatment ( $>2$  mg; Müller et al., 2017; Fernandez et al., 2017) typically required combining material from multiple (3–5) specimens grown under the same temperature conditions. To test potential inter-specimen differences, results were tracked per individual specimen for the  $1.1 \pm 0.2^{\circ}\text{C}$  and  $18 \pm 0.3^{\circ}\text{C}$  treatments (see **Table 1**).

### 2.2 Clumped isotope analysis

The clumped isotope composition of 278 aliquots of shell aragonite were analyzed over two 6-month periods (March – August 2020; May – November 2021) on two Thermo isotope ratio mass spectrometers (one MAT253 and one MAT253 plus) coupled to Kiel IV carbonate preparation devices (see **S1**). After correcting for variability in the pressure baseline (He et al., 2012), clumped isotope results were processed relative to the I-CDES through an empirical transfer function (ETF) based on measurements of ETH standards (ETH-1, -2 and -3) and their

accepted I-CDES values (Bernasconi et al., 2021). Isotopic values were calculated using the latest IUPAC values (Brand et al., 2010; Daëron et al., 2016). No AFF was applied after I-CDES standardization because the carbonate standards used for the ETF undergo the same acid reaction as the samples (Bernasconi et al., 2021). Long-term accuracy and reproducibility of  $\Delta_{47}$  results were assessed based on repeated measurements of the IAEA-C2 monitoring standard ( $\Delta_{47\_IAEA}$  on MAT253 plus:  $0.6382 \pm 0.026\text{‰}$ ;  $\Delta_{47\_IAEA}$  on MAT253:  $0.6445 \pm 0.046\text{‰}$ ;  $1\sigma$ ). Results were indistinguishable from the accepted value for IAEA-C2 ( $0.6409 \pm 0.003\text{‰}$ ; 95% CL; Bernasconi et al., 2021). Full results of all sample aliquots and standards used to standardize the results are provided in **S2**.

### 2.3 Data compilation

The *Arctica islandica* dataset was augmented with literature  $\Delta_{47}$  values of aragonites with known calcification temperatures (see **S3**). The dataset includes samples from mollusks (Bernasconi et al., 2018; aragonitic *Megapitaria aurantiaca* samples in Caldarescu et al., 2021; this study), foraminifera (Piasecki et al., 2019), travertines (Kele et al., 2015; Bernasconi et al., 2018), cave deposits (Breitenbach et al., 2018), lab-grown aragonites (Kluge et al., 2015) and heated aragonites (Müller et al., 2017). Data from several older studies (e.g. Ghosh et al., 2006; 2007; Tripathi et al., 2010; Wacker et al., 2013; 2014; Zhang et al., 2018; Zhai et al., 2019; Dong et al., 2021) were not included because they were not corrected for the pressure baseline (He et al., 2012; Bernasconi et al., 2013), could not be transferred into the standard reference frame (Dennis et al., 2011), lacked the standardization required to bring  $\Delta_{47}$  values into the I-CDES scale (Bernasconi et al., 2021) or because the aragonite was precipitated out of equilibrium (e.g. Kimball et al., 2015; Chen et al., 2019; **S1**; **S3**). Clumped isotope data from the literature was brought to the I-CDES reference frame using the multi-linear correction proposed in Appendix A of Bernasconi et al. (2021) using values of carbonate standards reported in the studies (see **S1**). Uncertainties on the formation temperatures of the non-temperature controlled datapoints from previous studies were generally in the order of  $1^\circ\text{C}$  ( $1\sigma$ ; see **S1**). The full dataset including  $\Delta_{47}$  values and temperatures with their uncertainties used

167 in this study is provided in **S4**. Unless stated otherwise, uncertainties are cited at the 95%  
168 confidence level.

169 All data processing for this study is done in R (R Core Team, 2022) and scripts are provided  
170 in **S5** and published on Github ([https://github.com/nielsidewinter/Aragonite\\_clumped](https://github.com/nielsidewinter/Aragonite_clumped)). Details  
171 on data processing are provided in **S1**. We compare our data with calibrations by Anderson et  
172 al. (2021) and Meinicke et al. (2020) as well as with temperature dependencies of aragonite  
173 and calcite clumped isotope compositions from *ab initio* modelling in Guo et al. (2009) brought  
174 into the I-CDES reference frame (see **S1**).



### 3. Results

#### 3.1 Clumped isotope values in *Arctica islandica*

Clumped isotope results from *A. islandica* are summarized in **Table 1** and **Figure 1**. There is no significant clumped isotope difference between specimens in the same temperature treatment ( $F(4,77) = 1.937$ ,  $p = 0.11$  for the 1°C specimens and  $F(3,63) = 0.377$ ,  $p = 0.77$  for the 18°C specimens; see **S6**). The number of measurements per specimen was large enough to exclude per-specimen  $\Delta_{47}$  differences outside the reproducibility standard deviation of the clumped isotope analyses (0.046‰; see **Table 1** and **S6**). Differences between all temperature treatments are statistically significant ( $P(3,274) = 15.68$ ,  $p < 0.01$ ), except for differences between the 15°C and 18°C temperature bin and the difference between 1°C and 3°C (95%CL; **S6**).

We investigated the  $\Delta_{47} - \frac{1}{T^2}$  relationship and how it varies along the temperature domain by performing linear regressions on increasingly large parts of our compilation. Note that the uncertainty on clumped isotope data compared to the range of temperatures of the lab-grown *A. islandica* leaves relatively high uncertainty on a clumped isotope-temperature regression through these results alone compared to the unified clumped isotope calibration (Anderson et al. 2021). We therefore do not advice using this and other regression equations in this section for calibrating clumped isotope results (see **Discussion**). Firstly, a statistically significant temperature relationship ( $\Delta_{47} - \frac{1}{T^2}$  slope  $> 0$ ; 95% CL) is found for  $\Delta_{47}$  exclusively from *Arctica islandica* samples:

$$\Delta_{47}(I - CDES) = 0.0280 \pm 0.0042 * \frac{10^6}{T^2} + 0.304 \pm 0.0524 \text{ (T in K, } \pm 1\sigma; \sigma_{res} = 0.047\text{‰)} \quad (1)$$

Secondly, including other aragonitic mollusk data (Caldarescu et al., 2021) yields a regression indistinguishable from the Anderson et al. (2021) unified clumped isotope calibration:

$$\Delta_{47}(I - CDES) = 0.0443 \pm 0.0024 * \frac{10^6}{T^2} + 0.097 \pm 0.0291 \text{ (T in K, } \pm 1\sigma; \sigma_{res} = 0.043\text{‰)} \quad (2)$$

197

### 198 3.2 Aragonite clumped isotope temperature dependence

199 When including clumped isotope values of other low-temperature (<30°C) aragonites in the  
200 compilation, the regression remains indistinguishable from the calibration of Anderson et al.  
201 (2021) and similar to Meinicke et al. (2020; 2021) and the Guo et al. (2009) theoretical  
202 temperature relationships (**Fig. 2B**):

$$\Delta_{47}(I - CDES) = 0.0451 \pm 0.0024 * \frac{10^6}{T^2} + 0.0871 \pm 0.0287 \text{ (T in K, } \pm 1\sigma; \sigma_{res} = 0.042\text{‰)} \quad (3)$$

203 Finally, we included higher temperature (>30°C) datapoints, such as the cave deposits of  
204 Breitenbach et al. (2018), travertine samples from Kele et al. (2015), precipitated aragonites  
205 from Kluge et al. (2015) and heated aragonites from Müller et al. (2017) in the linear regression.  
206 This decreases the slope and increases the intercept (see **Fig. 2**):

$$\Delta_{47}(I - CDES) = 0.0403 \pm 0.0005 * \frac{10^6}{T^2} + 0.1435 \pm 0.0485 \text{ (T in K, } \pm 1\sigma; \sigma_{res} = 0.040\text{‰)} \quad (4)$$

207 The formation temperatures of our *A. islandica* data on the very cold end of the calibration  
208 domain are significantly underestimated by Anderson et al. (2021;  $\Delta\Delta_{47} = +0.009 \pm 0.007\text{‰}$ ;  
209  $-2.71 \pm 2.03^\circ\text{C}$ ; **Fig. 3**;  $\Delta\Delta_{47}$  = offset between data and calibration). The theoretical aragonite  
210 clumped isotope-temperature relationship (Guo et al., 2009) severely overestimates *A.*  
211 *islandica* temperatures ( $-0.016 \pm 0.007\text{‰}$ ;  $+4.35 \pm 1.88^\circ\text{C}$ ; **Fig. 3**). Contrarily, the Meinicke et  
212 al. (2020; 2021) calibration ( $\Delta\Delta_{47} = +0.004 \pm 0.007\text{‰}$ ;  $-1.17 \pm 2.00^\circ\text{C}$ ; **Fig. 3**) and the  
213 theoretical calcite temperature relationship (Guo et al., 2009;  $\Delta\Delta_{47} = +0.002 \pm 0.007\text{‰}$ ;  $-0.47$   
214  $\pm 1.98^\circ\text{C}$ ; **Fig. 4**) do not significantly over- or underestimate the formation temperature of our  
215 *A. islandica* shells.

## 216 4. Discussion

### 217 4.1 Isotope ordering in aragonitic mollusks

Clumped isotope values of our temperature-controlled *A. islandica* samples consistently plot on a  $\Delta_{47} \sim \frac{1}{T^2}$  linear relationship with other low-temperature aragonite datapoints (**Fig. 1 and 2**; see **section 4.2**). The absence of a consistent offset between *A. islandica* datapoints and other aragonites (mean  $\Delta_{47}$  difference of  $+0.003 \pm 0.004\text{‰}$ , see **Fig. 2** and **S8**) and agreement between the linear  $\Delta_{47} \sim \frac{1}{T^2}$  dependence of the aragonitic mollusk data in this study and the regression through the complete low-temperature aragonite dataset (**Fig. 1** and **section 3.1**) strongly supports a common temperature dependence for all aragonites in this study, biogenic or inorganic, and argues against disequilibrium fractionation in aragonite precipitated inorganically or vital effect in bivalves or foraminifera (see **section 3.1**; **Fig. 1 and 2**). Our highly temperature-controlled growth experiments uniquely allow us to exclude variability in the growth environment between specimens from the same growth treatment as a driver of shell composition. Strong similarity of  $\Delta_{47}$  values between individual *A. islandica* specimens grown at the same temperature thus rules out specimen-specific vital effects on the clumped isotope composition aragonitic bivalve shells outside the uncertainty of our measurements (see **section 3.1**; **Fig. 1**, **Table 1** and **S6**). These findings corroborate measurements in calcitic mollusks showing that clumped isotope values in mollusk carbonates adhere to the same temperature relationship as other carbonates precipitated in equilibrium (except for juvenile oyster shells; Huyghe et al., 2022). Clumped isotope analyses in (fossil) mollusk shells thus provide an independent temperature proxy, allowing paleoclimatologists to disentangle the effects of variability in temperature and the hydrological cycle (as measured in  $\delta^{18}\text{O}_w$ ) throughout geological history down to the seasonal timescale (e.g. Caldarescu et al., 2021; de Winter et al., 2021; Letulle et al., 2022).

#### 4.2 Mineral-specific acid fractionation factor

Residuals of aragonite clumped isotope data around the low-temperature ( $<30^\circ\text{C}$ ) York regression ( $0.042\text{‰}$ ;  $1\sigma$ ; see **section 3.1** and **Fig. 2**) are predominantly explained by analytical uncertainty on  $\Delta_{47}$  measurements (external precision of  $0.026\text{‰}$  and  $0.046\text{‰}$  on the 253Plus and the MAT253 mass spectrometers;  $1\sigma$ ; see **section 2.2**). Uncertainty on formation

temperatures in the low-temperature dataset ( $\pm 0.8^\circ\text{C}$ ;  $1\sigma$ ; see **S4**) would add an additional uncertainty of  $0.0024\text{‰}$  ( $1\sigma$ ) if applied to the weighted average formation temperature of all low-temperature ( $<30^\circ\text{C}$ ) data points ( $22.0^\circ\text{C}$ ; see **S4**). Outside these uncertainties in the compilation data, there is little uncertainty on the temperature relationship in the low-temperature domain ( $<30^\circ\text{C}$ ; see **section 3.2**; **Fig. 2**). If clumped isotope fractionation during acid digestion was indeed different between aragonite and calcite (as suggested in Müller et al., 2017; Petersen et al., 2019) the difference in AFF would be indicated by the difference between our aragonite dataset and the previous calcite-based calibrations (Meinicke et al., 2020; 2021). The close similarity between our *A. islandica* data and the calcite calibration ( $\Delta\Delta_{47} = 0.004 \pm 0.007\text{‰}$ ; **Fig. 3**; **S7**) leaves little room for the  $0.007\text{‰}$  and  $0.025\text{‰}$  difference in AFF reported in Petersen et al. (2019) and Müller et al. (2017), respectively. We therefore conclude that the calcite AFF in Petersen et al. (2019), which are included in the I-CDES reference scale (Bernasconi et al., 2021) can be used for aragonite samples.

#### 4.3 Non-linear temperature dependence of clumped isotopes in aragonites

Current clumped isotope calibrations (Meinicke et al., 2020; 2021; Anderson et al., 2021) show subtle differences in the low-temperature end of the calibration ( $<30^\circ\text{C}$ ) that would result in  $\sim 1.5^\circ\text{C}$  colder temperatures when applying Anderson et al. (2021) compared to Meinicke et al. (2020). In addition, the cold-water ( $<30^\circ\text{C}$ ) carbonate based Meinicke et al. (2020) calibration more closely resembles the modelled temperature relationship for calcites in Guo et al. (2009). Including high-temperature ( $>30^\circ\text{C}$ ) data in our linear regression leads to overestimation of the temperature of warmer ( $>18^\circ\text{C}$ ) datapoints ( $\Delta\Delta_{47}$  of  $-0.005 \pm 0.006\text{‰}$ , or  $+1.8^{+2.1}_{-2.0}^\circ\text{C}$  for data precipitated at  $30^\circ\text{C}$ ), while underestimating colder datapoints ( $\Delta\Delta_{47}$  of  $+0.009 \pm 0.008\text{‰}$ , or  $-2.0^{+2.0}_{-2.0}^\circ\text{C}$  for data precipitated at  $0^\circ\text{C}$ ; **Fig. 2**; **S7**). Point-by-point offsets of all data from the calibration lines are provided in **S8**.

This difference between  $\Delta_{47} - \frac{1}{T^2}$  regressions through the low-temperature ( $<30^\circ\text{C}$ ) and the full dataset (see **section 3.2**; **Fig 2**) likely highlights non-linear behavior of the  $\Delta_{47} - \frac{1}{T^2}$  relationship

in aragonites. In fact, previous studies based on both clumped isotope analyses and *ab initio* modelling have suggested a non-linear  $\Delta_{47}-\frac{1}{T^2}$  relationship to be a better fit for both calcites (Guo et al., 2009; Jautzy et al., 2020) and dolomites (Guo et al., 2009; Müller et al., 2019) precipitated on a large range of known temperatures. Non-linear behavior is also observed in the Anderson et al. (2021) dataset, where  $\Delta_{47}$  values of calcites precipitated between 100°C and 1000°C are underestimated by the linear relationship, while the hottest datapoints (calcites heated to 1100°C) fall on the linear regression, mimicking the reduced  $\Delta_{47}-\frac{1}{T^2}$  slope at the high temperature end of the polynomial regressions through calcite and dolomite data (Guo et al., 2009; Jautzy et al., 2020; Müller et al., 2019). A linear  $\Delta_{47}-\frac{1}{T^2}$  relationship through a calibration dataset with a large temperature range will thus overestimate temperatures for samples with  $\Delta_{47}$  values between 0.2‰ and 0.4‰ (temperatures of 100°C–1000°C; see residuals in Anderson et al., 2021) and underestimate temperatures of cold (<30°C) samples, as confirmed by regressions through our low-temperature datapoints (see **Fig. 2-3** and **section 4.4**). Therefore, more high-temperature aragonite datapoints are needed to constrain the clumped isotope-temperature relationship for temperatures >100°C.

#### 4.4 Calibrating the clumped isotope-temperature relationship in cold (<30°C) carbonates

Our lab-grown *A. islandica* shells offer more control on formation temperature than naturally grown carbonates precipitated under variable temperatures. Ideally, the temperature of these natural samples is monitored so an average temperature can be calculated for the targeted growth period (e.g. Kele et al., 2015; de Winter et al., 2020; 2021b; Huyghe et al., 2021). However, formation temperatures are often indirectly estimated through other proxies (e.g.  $\delta^{18}\text{O}_c$ ) and/or estimates of the living environment (e.g. water depth) of the carbonate producer, accumulating uncertainty (e.g. Peral et al., 2018; Piasecki et al., 2018; Meinicke et al., 2020). These caveats obscure the full uncertainty of the formation temperatures of natural carbonates as well as the effect of this unknown uncertainty on the calibrations. Considering the methods by which the “known” temperatures of natural carbonates are estimated in previous studies,

part of the ~1.5°C temperature offset between Anderson et al. (2021) and Meinicke et al. (2020; 2021; see **Fig. 3**) and the  $2.7 \pm 2.0^\circ\text{C}$  offset between Anderson et al. (2021) and our *A. islandica* data might be caused by uncertainty on the formation temperatures of the calibration dataset. However, our highly temperature-controlled *A. islandica* datapoints reveal that, despite uncertainty on formation temperature, the Meinicke et al. (2021) calibration locally approximates the non-linear  $\Delta_{47} - \frac{1}{T^2}$  relationship in the cold temperature domain with higher accuracy than the Anderson et al. (2021) calibration (**Fig. 1, Fig. 3; S8**). The non-linear theoretical calcite temperature dependence by Guo et al. (2009) also fits well with the data. Precisely temperature-controlled carbonates thus better constrain the slope of the  $\Delta_{47} - \frac{1}{T^2}$  relationship for cold carbonates (improving calibration accuracy) while reducing the uncertainty on the calibration (improving calibration precision).

The ~1.5°C difference in reconstructed temperature between the calibrations in the low temperature range (<30°C) may seem trivial and requires the complete *A. islandica* dataset (N = 278; see **Fig. 4**) to resolve. However, in paleoclimate reconstructions (e.g. Petersen et al., 2016; de Winter et al., 2017; 2021a; Vickers et al., 2020b; Agterhuis et al., 2021; Meckler et al., 2022), this temperature offset may have significant consequences. A ~1.5°C cold bias in temperature reconstructions may lead to a significant underestimation of climate sensitivity to CO<sub>2</sub> forcing, biasing the physical science basis for informing policymakers about future climate change (e.g. Dennis et al., 2013; Modestou et al., 2020; Westerhold et al., 2020; Tierney et al., 2020; IPCC, 2021). Accurate clumped isotope-based temperature reconstructions therefore require calibration datasets with precisely constrained formation temperatures tailored to the temperature range of the samples.

## Acknowledgements

The authors would like to thank Nele Meckler and Stefano Bernasconi for their thoughtful review of the manuscript, and editor Angelicque White for moderating the review process.

Thanks to Arnold van Dijk and Desmond Eefting for their technical assistance in the UU clumped isotope lab. This work is part of the UNBIAS project jointly funded by a Flemish Research Foundation (FWO; 12ZB220N) post-doctoral fellowship (NJW) and a MSCA Individual Fellowship (H2020-MSCA-IF-2018; 843011 – UNBIAS; awarded to NJW). BG is supported by an UU-NIOZ collaboration grant.

## **Open Research**

Supplementary materials are deposited on the open-source repository Zenodo and can be accessed through the following link: <https://doi.org/10.5281/zenodo.6524705>. R scripts are uploaded on GitHub ([https://github.com/nielsidewinter/Aragonite\\_clumped](https://github.com/nielsidewinter/Aragonite_clumped)) and archived in Zenodo (<https://doi.org/10.5281/zenodo.6560188>).

## 334 References

- 335 Agterhuis, T., Ziegler, M., de Winter, N. J., and Lourens, L. J.: Warm deep-sea temperatures across  
 336 Eocene Thermal Maximum 2 from clumped isotope thermometry, *Commun Earth Environ*, 3, 1–9,  
 337 <https://doi.org/10.1038/s43247-022-00350-8>, 2022.
- 338 Anderson, N. T., Kelson, J. R., Kele, S., Daëron, M., Bonifacie, M., Horita, J., Mackey, T. J., John, C. M.,  
 339 Kluge, T., Petschnig, P., Jost, A. B., Huntington, K. W., Bernasconi, S. M., and Bergmann, K. D.: A  
 340 Unified Clumped Isotope Thermometer Calibration (0.5–1,100°C) Using Carbonate-Based  
 341 Standardization, 48, e2020GL092069, <https://doi.org/10.1029/2020GL092069>, 2021.
- 342 Bajnai, D., Guo, W., Spötl, C., Coplen, T. B., Methner, K., Löffler, N., Krsnik, E., Gischler, E., Hansen,  
 343 M., Henkel, D., Price, G. D., Raddatz, J., Scholz, D., and Fiebig, J.: Dual clumped isotope  
 344 thermometry resolves kinetic biases in carbonate formation temperatures, *Nat Commun*, 11,  
 345 4005, <https://doi.org/10.1038/s41467-020-17501-0>, 2020.
- 346 Bernasconi, S. M., Hu, B., Wacker, U., Fiebig, J., Breitenbach, S. F., and Rutz, T.: Background effects on  
 347 Faraday collectors in gas-source mass spectrometry and implications for clumped isotope  
 348 measurements, 27, 603–612, 2013.
- 349 Bernasconi, S. M., Müller, I. A., Bergmann, K. D., Breitenbach, S. F., Fernandez, A., Hodell, D. A., Jaggi,  
 350 M., Meckler, A. N., Millan, I., and Ziegler, M.: Reducing uncertainties in carbonate clumped  
 351 isotope analysis through consistent carbonate-based standardization, 19, 2895–2914, 2018.
- 352 Bernasconi, S. M., Daëron, M., Bergmann, K. D., Bonifacie, M., Meckler, A. N., Affek, H. P., Anderson,  
 353 N., Bajnai, D., Barkan, E., Beverly, E., Blamart, D., Burgener, L., Calmels, D., Chaduteau, C., Clog,  
 354 M., Davidheiser-Kroll, B., Davies, A., Dux, F., Eiler, J., Elliott, B., Fetrow, A. C., Fiebig, J., Goldberg,  
 355 S., Hermoso, M., Huntington, K. W., Hyland, E., Ingalls, M., Jaggi, M., John, C. M., Jost, A. B., Katz,  
 356 S., Kelson, J., Kluge, T., Kocken, I. J., Laskar, A., Leutert, T. J., Liang, D., Lucarelli, J., Mackey, T. J.,  
 357 Mangenot, X., Meinicke, N., Modestou, S. E., Müller, I. A., Murray, S., Neary, A., Packard, N.,  
 358 Passey, B. H., Pelletier, E., Petersen, S., Piasecki, A., Schauer, A., Snell, K. E., Swart, P. K., Tripathi, A.,  
 359 Upadhyay, D., Vennemann, T., Winkelstern, I., Yarian, D., Yoshida, N., Zhang, N., and Ziegler, M.:  
 360 InterCarb: A Community Effort to Improve Interlaboratory Standardization of the Carbonate  
 361 Clumped Isotope Thermometer Using Carbonate Standards, 22, e2020GC009588,  
 362 <https://doi.org/10.1029/2020GC009588>, 2021.
- 363 Breitenbach, S. F. M., Mlenek-Vautravers, M. J., Grauel, A.-L., Lo, L., Bernasconi, S. M., Müller, I. A.,  
 364 Rolfe, J., Gázquez, F., Greaves, M., and Hodell, D. A.: Coupled Mg/Ca and clumped isotope  
 365 analyses of foraminifera provide consistent water temperatures, *Geochimica et Cosmochimica*  
 366 *Acta*, 236, 283–296, <https://doi.org/10.1016/j.gca.2018.03.010>, 2018.
- 367 Buchardt, B. and Simonarson, L. A.: Isotope palaeotemperatures from the Tjörnes beds in Iceland:  
 368 evidence of Pliocene cooling, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 189, 71–95,  
 369 [https://doi.org/10.1016/S0031-0182\(02\)00594-1](https://doi.org/10.1016/S0031-0182(02)00594-1), 2003.
- 370 Butler, P. G., Wanamaker, A. D., Scourse, J. D., Richardson, C. A., and Reynolds, D. J.: Variability of  
 371 marine climate on the North Icelandic Shelf in a 1357-year proxy archive based on growth  
 372 increments in the bivalve *Arctica islandica*, 373, 141–151, 2013.
- 373 Caldarescu, D. E., Sadatzki, H., Andersson, C., Schäfer, P., Fortunato, H., and Meckler, A. N.: Clumped  
 374 isotope thermometry in bivalve shells: A tool for reconstructing seasonal upwelling, *Geochimica*  
 375 *et Cosmochimica Acta*, 294, 174–191, <https://doi.org/10.1016/j.gca.2020.11.019>, 2021.
- 376 Chen, S., Ryb, U., Piasecki, A. M., Lloyd, M. K., Baker, M. B., and Eiler, J. M.: Mechanism of solid-state  
 377 clumped isotope reordering in carbonate minerals from aragonite heating experiments,  
 378 *Geochimica et Cosmochimica Acta*, 258, 156–173, <https://doi.org/10.1016/j.gca.2019.05.018>,  
 379 2019.



- Daëron, M., Blamart, D., Peral, M., and Affek, H. P.: Absolute isotopic abundance ratios and the accuracy of  $\Delta 47$  measurements, 442, 83–96, 2016.
- Daëron, M., Drysdale, R. N., Peral, M., Huyghe, D., Blamart, D., Coplen, T. B., Lartaud, F., and Zanchetta, G.: Most Earth-surface calcites precipitate out of isotopic equilibrium, 10, 429, <https://doi.org/10.1038/s41467-019-08336-5>, 2019.
- De Winter, N., Vellekoop, J., Vorrsselmans, R., Golreihan, A., Soete, J., Petersen, S., Meyer, K., Casadío, S., Speijer, R., and Claeys, P.: An assessment of latest Cretaceous Pycnodonte vesicularis (Lamarck, 1806) shells as records for palaeoseasonality: A multi-proxy investigation, *Climate of the Past Discussions*, 2017, 1–36, 2017.
- Deming, W. E.: Statistical adjustment of data., 1943.
- Dennis, K. J. and Schrag, D. P.: Clumped isotope thermometry of carbonatites as an indicator of diagenetic alteration, *Geochimica et Cosmochimica Acta*, 74, 4110–4122, <https://doi.org/10.1016/j.gca.2010.04.005>, 2010.
- Dennis, K. J., Affek, H. P., Passey, B. H., Schrag, D. P., and Eiler, J. M.: Defining an absolute reference frame for ‘clumped’ isotope studies of CO<sub>2</sub>, *Geochimica et Cosmochimica Acta*, 75, 7117–7131, <https://doi.org/10.1016/j.gca.2011.09.025>, 2011.
- Dennis, K. J., Cochran, J. K., Landman, N. H., and Schrag, D. P.: The climate of the Late Cretaceous: New insights from the application of the carbonate clumped isotope thermometer to Western Interior Seaway macrofossil, *Earth and Planetary Science Letters*, 362, 51–65, <https://doi.org/10.1016/j.epsl.2012.11.036>, 2013.
- Dong, J., Eiler, J., An, Z., Li, X., Liu, W., and Hu, J.: Clumped isotopic compositions of cultured and natural land-snail shells and their implications, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 577, 110530, <https://doi.org/10.1016/j.palaeo.2021.110530>, 2021.
- Eiler, J. M.: “Clumped-isotope” geochemistry—The study of naturally-occurring, multiply-substituted isotopologues, *Earth and Planetary Science Letters*, 262, 309–327, <https://doi.org/10.1016/j.epsl.2007.08.020>, 2007.
- EPSTEIN, S., BUCHSBAUM, R., LOWENSTAM, H. A., and UREY, H. C.: REVISED CARBONATE-WATER ISOTOPIC TEMPERATURE SCALE, *GSA Bulletin*, 64, 1315–1326, [https://doi.org/10.1130/0016-7606\(1953\)64\[1315:RCITS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1953)64[1315:RCITS]2.0.CO;2), 1953.
- Fernandez, A., Müller, I. A., Rodríguez-Sanz, L., van Dijk, J., Looser, N., and Bernasconi, S. M.: A reassessment of the precision of carbonate clumped isotope measurements: implications for calibrations and paleoclimate reconstructions, 18, 4375–4386, 2017.
- Fiebig, J., Daëron, M., Bernecker, M., Guo, W., Schneider, G., Boch, R., Bernasconi, S. M., Jautzy, J., and Dietzel, M.: Calibration of the dual clumped isotope thermometer for carbonates, *Geochimica et Cosmochimica Acta*, <https://doi.org/10.1016/j.gca.2021.07.012>, 2021.
- Frisia, S., Borsato, A., Fairchild, I. J., and McDermott, F.: Calcite Fabrics, Growth Mechanisms, and Environments of Formation in Speleothems from the Italian Alps and Southwestern Ireland, *Journal of Sedimentary Research*, 70, 1183–1196, <https://doi.org/10.1306/022900701183>, 2000.
- Ghosh, P., Adkins, J., Affek, H., Balta, B., Guo, W., Schauble, E. A., Schrag, D., and Eiler, J. M.: 13C–18O bonds in carbonate minerals: A new kind of paleothermometer, *Geochimica et Cosmochimica Acta*, 70, 1439–1456, <https://doi.org/10.1016/j.gca.2005.11.014>, 2006.
- Ghosh, P., Eiler, J., Campana, S. E., and Feeney, R. F.: Calibration of the carbonate ‘clumped isotope’ paleothermometer for otoliths, *Geochimica et Cosmochimica Acta*, 71, 2736–2744, <https://doi.org/10.1016/j.gca.2007.03.015>, 2007.
- Goodwin, D. H., Flessa, K. W., Schöne, B. R., and Dettman, D. L.: Cross-calibration of daily growth increments, stable isotope variation, and temperature in the Gulf of California bivalve mollusk *Chione cortezi*: implications for paleoenvironmental analysis, 16, 387–398, 2001.

- Goodwin, D. H., Schöne, B. R., and Dettman, D. L.: Resolution and Fidelity of Oxygen Isotopes as Paleotemperature Proxies in Bivalve Mollusk Shells: Models and Observations, *PALAIOS*, 18, 110–125, [https://doi.org/10.1669/0883-1351\(2003\)18<110:RAFOOI>2.0.CO;2](https://doi.org/10.1669/0883-1351(2003)18<110:RAFOOI>2.0.CO;2), 2003.
- Guo, W.: Kinetic clumped isotope fractionation in the DIC-H<sub>2</sub>O-CO<sub>2</sub> system: Patterns, controls, and implications, *Geochimica et Cosmochimica Acta*, 268, 230–257, <https://doi.org/10.1016/j.gca.2019.07.055>, 2020.
- Guo, W., Mosenfelder, J. L., Goddard, W. A., and Eiler, J. M.: Isotopic fractionations associated with phosphoric acid digestion of carbonate minerals: Insights from first-principles theoretical modeling and clumped isotope measurements, *Geochimica et Cosmochimica Acta*, 73, 7203–7225, <https://doi.org/10.1016/j.gca.2009.05.071>, 2009.
- Hansen, H. J.: Test structure and evolution in the Foraminifera, 12, 173–182, <https://doi.org/10.1111/let.1979.12.2.173>, 1979.
- He, B., Olack, G. A., and Colman, A. S.: Pressure baseline correction and high-precision CO<sub>2</sub> clumped-isotope ( $\Delta 47$ ) measurements in bellows and micro-volume modes, 26, 2837–2853, 2012.
- Henkes, G. A., Passey, B. H., Wanamaker, A. D., Grossman, E. L., Ambrose, W. G., and Carroll, M. L.: Carbonate clumped isotope compositions of modern marine mollusk and brachiopod shells, *Geochimica et Cosmochimica Acta*, 106, 307–325, <https://doi.org/10.1016/j.gca.2012.12.020>, 2013.
- Henkes, G. A., Passey, B. H., Grossman, E. L., Shenton, B. J., Yancey, T. E., and Pérez-Huerta, A.: Temperature evolution and the oxygen isotope composition of Phanerozoic oceans from carbonate clumped isotope thermometry, *Earth and Planetary Science Letters*, 490, 40–50, <https://doi.org/10.1016/j.epsl.2018.02.001>, 2018.
- Huyghe, D., Daëron, M., de Rafelis, M., Blamart, D., Sébilo, M., Paulet, Y.-M., and Lartaud, F.: Clumped isotopes in modern marine bivalves, *Geochimica et Cosmochimica Acta*, 316, 41–58, <https://doi.org/10.1016/j.gca.2021.09.019>, 2022.
- Jautzy, J. J., Savard, M. M., Dhillon, R. S., Bernasconi, S. M., and Smirnov, A.: Clumped isotope temperature calibration for calcite: Bridging theory and experimentation, 14, 36–41, 2020.
- Kele, S., Breitenbach, S. F., Capezzuoli, E., Meckler, A. N., Ziegler, M., Millan, I. M., Kluge, T., Deák, J., Hanselmann, K., and John, C. M.: Temperature dependence of oxygen-and clumped isotope fractionation in carbonates: a study of travertines and tufas in the 6–95 °C temperature range, 168, 172–192, 2015.
- Kennedy, W. J., Taylor, J. D., and Hall, A.: Environmental and Biological Controls on Bivalve Shell Mineralogy, 44, 499–530, <https://doi.org/10.1111/j.1469-185X.1969.tb00610.x>, 1969.
- Kim, S.-T. and O’Neil, J. R.: Equilibrium and nonequilibrium oxygen isotope effects in synthetic carbonates, *Geochimica et Cosmochimica Acta*, 61, 3461–3475, [https://doi.org/10.1016/S0016-7037\(97\)00169-5](https://doi.org/10.1016/S0016-7037(97)00169-5), 1997.
- Kimball, J., Eagle, R., and Dunbar, R.: Carbonate “clumped” isotope signatures in aragonitic scleractinian and calcitic gorgonian deep-sea corals, 13, 6487–6505, <https://doi.org/10.5194/bg-13-6487-2016>, 2016.
- Kirchner, J.W. Data Analysis Toolkit #12: Weighted averages and their uncertainty: [http://seismo.berkeley.edu/~kirchner/Toolkits/Toolkit\\_12.pdf](http://seismo.berkeley.edu/~kirchner/Toolkits/Toolkit_12.pdf), last access: 3 August 2022.
- Kluge, T., John, C. M., Jourdan, A.-L., Davis, S., and Crawshaw, J.: Laboratory calibration of the calcium carbonate clumped isotope thermometer in the 25–250 °C temperature range, *Geochimica et Cosmochimica Acta*, 157, 213–227, <https://doi.org/10.1016/j.gca.2015.02.028>, 2015.
- Knutti, R., Rugenstein, M. A. A., and Hegerl, G. C.: Beyond equilibrium climate sensitivity, *Nature Geosci*, 10, 727–736, <https://doi.org/10.1038/ngeo3017>, 2017.

- Kocken, I. J., Müller, I. A., and Ziegler, M.: Optimizing the Use of Carbonate Standards to Minimize Uncertainties in Clumped Isotope Data, 20, 5565–5577, <https://doi.org/10.1029/2019GC008545>, 2019.
- Letulle, T., Suan, G., Daëron, M., Rogov, M., Lécuyer, C., Vinçon-Laugier, A., Reynard, B., Montagnac, G., Lutikov, O., and Schlögl, J.: Clumped isotope evidence for Early Jurassic extreme polar warmth and high climate sensitivity, 18, 435–448, <https://doi.org/10.5194/cp-18-435-2022>, 2022.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, Ö., Yu, R., and Zhou, B. (Eds.): Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2021.
- Meckler, A. N., Ziegler, M., Millán, M. I., Breitenbach, S. F., and Bernasconi, S. M.: Long-term performance of the Kiel carbonate device with a new correction scheme for clumped isotope measurements, 28, 1705–1715, 2014.
- Meinicke, N., Ho, S. L., Hannisdal, B., Nürnberg, D., Tripathi, A., Schiebel, R., and Meckler, A. N.: A robust calibration of the clumped isotopes to temperature relationship for foraminifers, *Geochimica et Cosmochimica Acta*, 270, 160–183, <https://doi.org/10.1016/j.gca.2019.11.022>, 2020.
- Meinicke, N., Reimi, M. A., Ravelo, A. C., and Meckler, A. N.: Coupled Mg/Ca and Clumped Isotope Measurements Indicate Lack of Substantial Mixed Layer Cooling in the Western Pacific Warm Pool During the Last ~5 Million Years, 36, e2020PA004115, <https://doi.org/10.1029/2020PA004115>, 2021.
- Modestou, S. E., Leutert, T. J., Fernandez, A., Lear, C. H., and Meckler, A. N.: Warm Middle Miocene Indian Ocean Bottom Water Temperatures: Comparison of Clumped Isotope and Mg/Ca-Based Estimates, 35, e2020PA003927, <https://doi.org/10.1029/2020PA003927>, 2020.
- Müller, I. A., Violay, M. E. S., Storck, J.-C., Fernandez, A., van Dijk, J., Madonna, C., and Bernasconi, S. M.: Clumped isotope fractionation during phosphoric acid digestion of carbonates at 70°C, *Chemical Geology*, 449, 1–14, <https://doi.org/10.1016/j.chemgeo.2016.11.030>, 2017.
- Müller, I. A., Rodriguez-Blanco, J. D., Storck, J.-C., do Nascimento, G. S., Bontognali, T. R. R., Vasconcelos, C., Benning, L. G., and Bernasconi, S. M.: Calibration of the oxygen and clumped isotope thermometers for (proto-)dolomite based on synthetic and natural carbonates, *Chemical Geology*, 525, 1–17, <https://doi.org/10.1016/j.chemgeo.2019.07.014>, 2019.
- Nooitgedacht, C. W., van der Lubbe, H. J. L., Ziegler, M., and Staudigel, P. T.: Internal Water Facilitates Thermal Resetting of Clumped Isotopes in Biogenic Aragonite, 22, e2021GC009730, <https://doi.org/10.1029/2021GC009730>, 2021.
- Peral, M., Daëron, M., Blamart, D., Bassinot, F., Dewilde, F., Smialkowski, N., Isguder, G., Bonnin, J., Jorissen, F., and Kissel, C.: Updated calibration of the clumped isotope thermometer in planktonic and benthic foraminifera, 239, 1–16, 2018.
- Petersen, S. V., Tabor, C. R., Lohmann, K. C., Poulsen, C. J., Meyer, K. W., Carpenter, S. J., Erickson, J. M., Matsunaga, K. K., Smith, S. Y., and Sheldon, N. D.: Temperature and salinity of the Late Cretaceous western interior seaway, 44, 903–906, 2016.
- Petersen, S. V., Defliese, W. F., Saenger, C., Daëron, M., Huntington, K. W., John, C. M., Kelson, J. R., Bernasconi, S. M., Colman, A. S., Kluge, T., Olack, G. A., Schauer, A. J., Bajnai, D., Bonifacie, M., Breitenbach, S. F. M., Fiebig, J., Fernandez, A. B., Henkes, G. A., Hodell, D., Katz, A., Kele, S., Lohmann, K. C., Passey, B. H., Peral, M. Y., Petrizzo, D. A., Rosenheim, B. E., Tripathi, A., Venturelli, R., Young, E. D., and Winkelstern, I. Z.: Effects of Improved  $\delta^{18}\text{O}$  Correction on Interlaboratory Agreement in Clumped Isotope Calibrations, Estimates of Mineral-Specific Offsets, and

520 Temperature Dependence of Acid Digestion Fractionation, 20, 3495–3519,  
 521 <https://doi.org/10.1029/2018GC008127>, 2019.

522 Piasecki, A., Bernasconi, S. M., Grauel, A.-L., Hannisdal, B., Ho, S. L., Leutert, T. J., Marchitto, T. M.,  
 523 Meinicke, N., Tisserand, A., and Meckler, N.: Application of Clumped Isotope Thermometry to  
 524 Benthic Foraminifera, 20, 2082–2090, <https://doi.org/10.1029/2018GC007961>, 2019.

525 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical  
 526 Computing, Vienna, Austria, 2022.

527 Rodríguez-Sanz, L., Bernasconi, S. M., Marino, G., Heslop, D., Müller, I. A., Fernandez, A., Grant, K. M.,  
 528 and Rohling, E. J.: Penultimate deglacial warming across the Mediterranean Sea revealed by  
 529 clumped isotopes in foraminifera, Sci Rep, 7, 16572, [https://doi.org/10.1038/s41598-017-16528-](https://doi.org/10.1038/s41598-017-16528-6)  
 530 [6](https://doi.org/10.1038/s41598-017-16528-6), 2017.

531 Schaefer, R., Trutschler, K., and Rumohr, H.: Biometric studies on the bivalves *Astarte elliptica*, *A.*  
 532 *borealis* and *A. montagui* in Kiel Bay (Western Baltic Sea), *Helgolander Meeresunters*, 39, 245–253,  
 533 <https://doi.org/10.1007/BF01992772>, 1985.

534 Schauble, E. A., Eiler, J. M., and Kitchen, N.: Measurement and significance of  $^{13}\text{C}^{18}\text{O}^{16}\text{O}$  in  
 535 thermodynamically equilibrated and environmental  $\text{CO}_2$ , 67, A419–A419, 2003.

536 Schauble, E. A., Ghosh, P., and Eiler, J. M.: Preferential formation of  $^{13}\text{C}$ – $^{18}\text{O}$  bonds in carbonate  
 537 minerals, estimated using first-principles lattice dynamics, *Geochimica et Cosmochimica Acta*, 70,  
 538 2510–2529, <https://doi.org/10.1016/j.gca.2006.02.011>, 2006.

539 Schöne, B. R. and Fiebig, J.: Seasonality in the North Sea during the Allerød and Late Medieval  
 540 Climate Optimum using bivalve sclerochronology, 98, 83–98, 2009.

541 Schöne, B. R., Fiebig, J., Pfeiffer, M., Gleß, R., Hickson, J., Johnson, A. L., Dreyer, W., and Oschmann,  
 542 W.: Climate records from a bivalved *Methuselah* (*Arctica islandica*, Mollusca; Iceland), 228, 130–  
 543 148, 2005.

544 Staudigel, P. T. and Swart, P. K.: Isotopic behavior during the aragonite-calcite transition: Implications  
 545 for sample preparation and proxy interpretation, *Chemical Geology*, 442, 130–138,  
 546 <https://doi.org/10.1016/j.chemgeo.2016.09.013>, 2016.

547 Sturm, P.: bfit: Best-Fit Straight Line, 2018.

548 Swart, P. K., Lu, C., Moore, E. W., Smith, M. E., Murray, S. T., and Staudigel, P. T.: A calibration  
 549 equation between  $\Delta 48$  values of carbonate and temperature, 35, e9147,  
 550 <https://doi.org/10.1002/rcm.9147>, 2021.

551 Tatebe, K. Combining Multiple Averaged Data Points And Their Errors:  
 552 <https://docplayer.net/33088897-Combining-multiple-averaged-data-points-and-their-errors.html>,  
 553 last access: 3 August 2022.

554 Taylor, J. D. and Reid, D. G.: Shell microstructure and mineralogy of the Littorinidae: ecological and  
 555 evolutionary significance, in: *Progress in Littorinid and Muricid Biology*, Dordrecht, 199–215,  
 556 [https://doi.org/10.1007/978-94-009-0563-4\\_16](https://doi.org/10.1007/978-94-009-0563-4_16), 1990.

557 Tierney, J. E., Poulsen, C. J., Montañez, I. P., Bhattacharya, T., Feng, R., Ford, H. L., Hönisch, B., Inglis,  
 558 G. N., Petersen, S. V., Sagoo, N., Tabor, C. R., Thirumalai, K., Zhu, J., Burls, N. J., Foster, G. L.,  
 559 Goddérís, Y., Huber, B. T., Ivany, L. C., Turner, S. K., Lunt, D. J., McElwain, J. C., Mills, B. J. W., Otto-  
 560 Bliesner, B. L., Ridgwell, A., and Zhang, Y. G.: Past climates inform our future, 370,  
 561 <https://doi.org/10.1126/science.aay3701>, 2020.

562 Tripathi, A. K., Eagle, R. A., Thiagarajan, N., Gagnon, A. C., Bauch, H., Halloran, P. R., and Eiler, J. M.:  
 563  $^{13}\text{C}$ – $^{18}\text{O}$  isotope signatures and ‘clumped isotope’ thermometry in foraminifera and coccoliths,  
 564 *Geochimica et Cosmochimica Acta*, 74, 5697–5717, <https://doi.org/10.1016/j.gca.2010.07.006>,  
 565 2010.

566 Urey, H. C.: The thermodynamic properties of isotopic substances - Google Scholar, 562–581, 1947.


- Vickers, M. L., Lengger, S. K., Bernasconi, S. M., Thibault, N., Schultz, B. P., Fernandez, A., Ullmann, C. V., McCormack, P., Bjerrum, C. J., Rasmussen, J. A., Hougård, I. W., and Korte, C.: Cold spells in the Nordic Seas during the early Eocene Greenhouse, *Nat Commun*, 11, 4713, <https://doi.org/10.1038/s41467-020-18558-7>, 2020a.
- Vickers, M. L., Fernandez, A., Hesselbo, S. P., Price, G. D., Bernasconi, S. M., Lode, S., Ullmann, C. V., Thibault, N., Hougård, I. W., and Korte, C.: Unravelling Middle to Late Jurassic palaeoceanographic and palaeoclimatic signals in the Hebrides Basin using belemnite clumped isotope thermometry, *Earth and Planetary Science Letters*, 546, 116401, <https://doi.org/10.1016/j.epsl.2020.116401>, 2020b.
- Wacker, U., Fiebig, J., and Schoene, B. R.: Clumped isotope analysis of carbonates: comparison of two different acid digestion techniques, 27, 1631–1642, 2013.
- Wacker, U., Fiebig, J., Tödter, J., Schöne, B. R., Bahr, A., Friedrich, O., Tütken, T., Gischler, E., and Joachimski, M. M.: Empirical calibration of the clumped isotope paleothermometer using calcites of various origins, 141, 127–144, 2014.
- Wang, Z., Schauble, E. A., and Eiler, J. M.: Equilibrium thermodynamics of multiply substituted isotopologues of molecular gases, *Geochimica et Cosmochimica Acta*, 68, 4779–4797, <https://doi.org/10.1016/j.gca.2004.05.039>, 2004.
- Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnett, J. S., Bohaty, S. M., De Vleeschouwer, D., and Florindo, F.: An astronomically dated record of Earth's climate and its predictability over the last 66 million years, 369, 1383–1387, 2020.
- de Winter, N. J., Vellekoop, J., Clark, A. J., Stassen, P., Speijer, R. P., and Claeys, P.: The giant marine gastropod *Campanile giganteum* (Lamarck, 1804) as a high-resolution archive of seasonality in the Eocene greenhouse world, 21, e2019GC008794, <https://doi.org/10.1029/2019GC008794>, 2020.
- de Winter, N. J., Müller, I. A., Kocken, I. J., Thibault, N., Ullmann, C. V., Farnsworth, A., Lunt, D. J., Claeys, P., and Ziegler, M.: Absolute seasonal temperature estimates from clumped isotopes in bivalve shells suggest warm and variable greenhouse climate, *Commun Earth Environ*, 2, 1–8, <https://doi.org/10.1038/s43247-021-00193-9>, 2021a.
- de Winter, N. J., Dämmer, L. K., Falkenroth, M., Reichart, G.-J., Moretti, S., Martínez-García, A., Höche, N., Schöne, B. R., Rodiouchkina, K., Goderis, S., Vanhaecke, F., van Leeuwen, S. M., and Ziegler, M.: Multi-isotopic and trace element evidence against different formation pathways for oyster microstructures, *Geochimica et Cosmochimica Acta*, 308, 326–352, <https://doi.org/10.1016/j.gca.2021.06.012>, 2021b.
- de Winter, Niels J., Witbaard, Rob, Kocken, Ilja J., Müller, Inigo A., Guo, Jingjing, Goudsmit, Barbara, & Ziegler, Martin. (2022). Supplementary Information for "Temperature dependence of clumped isotopes ( $\Delta 47$ ) in aragonite". <https://doi.org/10.5281/zenodo.6524706>
- de Winter, N. J.: R Scripts belonging to: Temperature dependence of clumped isotopes ( $\Delta 47$ ) in aragonite, GitHub, [https://github.com/nielsjdewinter/Aragonite\\_clumped](https://github.com/nielsjdewinter/Aragonite_clumped), 2022. Witbaard, R., Duineveld, G. C. A., and De Wilde, P.: A long-term growth record derived from *Arctica islandica* (Mollusca, Bivalvia) from the Fladen Ground (northern North Sea), 77, 801–816, 1997.
- Witbaard, R., Franken, R., and Visser, B.: Growth of juvenile *Arctica islandica* under experimental conditions, *Helgoländer Meeresunters.*, 51, 417, <https://doi.org/10.1007/BF02908724>, 1998.
- Zhai, J., Wang, X., Qin, B., Cui, L., Zhang, S., and Ding, Z.: Clumped isotopes in land snail shells over China: Towards establishing a biogenic carbonate paleothermometer, *Geochimica et Cosmochimica Acta*, 257, 68–79, <https://doi.org/10.1016/j.gca.2019.04.028>, 2019.
- Zhang, N., Yamada, K., Kano, A., Matsumoto, R., and Yoshida, N.: Equilibrated clumped isotope signatures of land-snail shells observed from laboratory culturing experiments and its environmental implications, *Chemical Geology*, 488, 189–199, <https://doi.org/10.1016/j.chemgeo.2018.05.001>, 2018.

**Table 1 *Arctica islandica* clumped isotope results compared to previous calibrations**


“mixed” = combined samples from multiple specimens, number codes (e.g. “29” or “6A”) = material from one individual. “Ais” = *Arctica islandica*. Significant  $\Delta_{47}$  and temperature offsets are labeled in bold. cc = calcite, ar = aragonite

Sample	Culturing temperature	$\Delta_{47}$ (I-CDES; $\pm$ 95%CL)	N	Offset from Anderson	Offset from Meinicke	Offset from Guo (cc)	Offset from Guo (ar)
<b>Ais1</b>	$1.1 \pm 0.2^\circ\text{C}$	mixed	$0.695 \pm 0.019\text{‰}$	34			
		3	$0.661 \pm 0.023\text{‰}$	13			
		29	$0.688 \pm 0.025\text{‰}$	15			
		6A	$0.684 \pm 0.028\text{‰}$	11			
		6B	$0.649 \pm 0.026\text{‰}$	9			
		TOTAL	$0.682 \pm 0.010\text{‰}$	82	$+0.008\text{‰}$ $-2.12^\circ\text{C}$	$+0.002\text{‰}$ $-0.61^\circ\text{C}$	$+0.003\text{‰}$ $-0.75^\circ\text{C}$ <b><math>-0.016\text{‰}</math></b> <b><math>+4.48^\circ\text{C}</math></b>
<b>Ais3</b>	$3.2 \pm 0.3^\circ\text{C}$	mixed	$0.667 \pm 0.010\text{‰}$	72	$+0.001\text{‰}$ $-0.27^\circ\text{C}$	$-0.005\text{‰}$ $+1.24^\circ\text{C}$	$-0.006\text{‰}$ $+1.57^\circ\text{C}$ <b><math>-0.023\text{‰}</math></b> <b><math>+6.80^\circ\text{C}</math></b>
<b>Ais15</b>	$15 \pm 0.4^\circ\text{C}$	mixed	$0.637 \pm 0.009\text{‰}$	57	<b><math>+0.013\text{‰}</math></b> <b><math>-3.63^\circ\text{C}</math></b>	$+0.008\text{‰}$ $-2.10^\circ\text{C}$	$+0.004\text{‰}$ $-0.99^\circ\text{C}$ <b><math>-0.013\text{‰}</math></b> <b><math>+4.25^\circ\text{C}</math></b>
<b>Ais18</b>	$18 \pm 0.3^\circ\text{C}$	mixed	$0.635 \pm 0.010\text{‰}$	39			
		67	$0.647 \pm 0.024\text{‰}$	9			
		89	$0.640 \pm 0.028\text{‰}$	8			
		111	$0.630 \pm 0.022\text{‰}$	11			
		TOTAL	$0.637 \pm 0.005\text{‰}$	67	<b><math>+0.021\text{‰}</math></b> <b><math>-6.63^\circ\text{C}</math></b>	<b><math>+0.016\text{‰}</math></b> <b><math>-5.10^\circ\text{C}</math></b>	<b><math>+0.012\text{‰}</math></b> <b><math>-3.99^\circ\text{C}</math></b> $-0.004\text{‰}$ $+1.25^\circ\text{C}$






**Figure 1: *Arctica islandica*  $\Delta_{47}$  results.** Clumped isotope results are aggregated by specimen or multi-specimen sample (round symbols; see **Table 1**). Vertical lines represent 95% CL and number indicate sample size. Data is color-coded per temperature treatment (1°C, 3°C, 15°C and 18°C), with bold error bars indicating 95% CL, pairs of letter labels (a and b) indicate statistically indistinguishable  $\Delta_{47}$  values ( $p < 0.05$ ). The grey error bar at 6°C highlights *A. islandica* data from Bernasconi et al. (2018; recalculated to I-CDES). Solid and dashed black lines show calibrations by Anderson et al. (2021) and Meinicke et al. (2020; projected on I-CDES scale; Meinicke et al., 2021), respectively. Grey solid and dashed lines represent, respectively, the theoretical calcite (“cc”) and aragonite (“ar”) temperature dependencies from Guo et al. (2009; projected on the I-CDES scale, see **S1**). The horizontal axis is scaled to  $\frac{10^6}{T^2}$ , with T in K, to show the assumed linear relationship with the clumped isotope value.



**Figure 2: Aragonite  $\Delta_{47}$  temperature dependence.** Clumped isotope data of aragonite samples plotted against formation temperature. **A.** All data plotted over the full temperature range (1°C–850°C). Individual datapoints, averages and uncertainty on temperature and  $\Delta_{47}$  values (95% CL) are color-coded by study. Symbols highlight different types of aragonite. The solid and dashed black lines show calibrations by Anderson et al. (2021) and Meinicke et al. (2020; 2021; plotted for temperatures <30°C). Grey solid and dashed lines represent, respectively, the theoretical calcite ("cc") and aragonite ("ar") temperature dependencies from Guo et al. (2009; projected on the I-CDES scale, see **section 2.5**). Colored dashed lines and shaded envelopes show York regressions through aragonite data and their 95% confidence envelopes, respectively. **B.** Shows a zoom-in of the plot in **A.** for the low-temperature domain (1–30°C). Note that the horizontal axis is scaled to  $\frac{10^6}{T^2}$ , with T in K, to show the assumed linear relationship with the clumped isotope value.





**Figure 3: Offset of *A. islandica* data from temperature regressions.** Shaded grey points show residual  $\Delta_{47}$  values relative to four clumped isotope temperature relationships (see horizontal axis). Black symbols with error bars (95% CL) show mean offsets of all *A. islandica* datapoints (grown at 1°C, 3°C, 6°C, 15°C and 18°C) from the calibrations. The vertical axis on the right shows the temperature offset relative to the weighted mean calcification temperature of the full *A. islandica* dataset (8.6°C; see **S4**) based on Anderson et al. (2021).