

```

---
title: "Ocean Alkalinity, Buffering and Biogeochemical Processes -
accompanying scripts"
author: "Karline Soetaert and Mathilde Hagens"
params:
  EVAL: no
output:
  pdf_document: default
  word_document: default
  html_document: default
---

```{r setup, include=FALSE}
knitr::opts_chunk$set(echo = FALSE)
```

# Preamble

These scripts were used for the calculations in the Middelburg et al (2020)
paper.

```{r, message=FALSE}
require(AquaEnv)
require(plot3D)
require(RColorBrewer)

darkcols <- brewer.pal(12,"Paired")
darkcols[11] <- grey(0.1)
palette(darkcols)

ColScheme <- ramp.col(col = c("#1F78B4", "#FFFFCC", "#FC9272") , n =
100)
```

# Environmental settings

The default species composition as in Hagens and Middelburg (2016) is used.

```{r}

Hagens and middelburg (2016):
SETTINGS <- list(SumCO2 = 2017e-6, SumBOH3 = 427.8e-6, SumH3PO4 = 0.5e-6,
 SumSiOH4 = 7.4e-6, SumH2S = 2.3e-9, SumH2SO4 = 27930e-6,
 SumHF = 67.6e-6, SumHNO3 = 5.2e-6, SumHNO2 = 0.1e-6,
 SumH2O = 55.49, SumNH4 =0.3e-6, S = 34.617, t = 18.252, p
= 0)

Lumpsum <- data.frame(lumpsum = unlist(SETTINGS[c("SumCO2", "SumBOH3",
"SumH3PO4", "SumSiOH4",
 "SumH2S", "SumH2SO4", "SumHF", "SumHNO3",
 "SumHNO2", "SumH2O", "SumNH4")]))

expression <- list(expression(sum("CO"[2])), expression(sum("B(OH)"[3])),
 expression(sum(paste("H"[3],"PO"[4]))),
expression(sum("Si(OH)"[4])),
 expression(sum(paste("H"[2],"S"))),
expression(sum(paste("H"[2],"SO"[4]))),
 expression(sum("HF")), expression(sum("HNO"[3])),
 expression(sum("HNO"[2])),
expression(sum(paste("H"[2],"O"))),
 expression(sum("NH"[4])))
Lumpsum$expression <- expression

Redfield ratios
NC <- 16/106
PC <- 1/106

pH.seq <- seq(1, 14, length.out = 100)

```

```

...

Chemical players

The various chemical species are inputted here, together with their
charges, their contributions to alkalinity, and the lump sum to which they
belong:

```{r}
Speciation <- data.frame(
  species = c("CO2", "HCO3", "CO3",
             "BOH3", "BOH4",
             "H3PO4", "H2PO4", "HPO4", "PO4",
             "SiOH4", "SiOOH3", "SiO2OH2",
             "H2S", "HS", "S2min",
             "H2SO4", "HSO4", "SO4",
             "HF", "F",
             "HNO3", "NO3",
             "HNO2", "NO2",
             "OH", "H2O", "H",
             "NH4", "NH3"),

             #SumCO2 SumBOH3 SumH3PO4 SumSiOH4 SumH2S SumH2SO4 SumHF
SumHNO3 SumHNO2 SumH2O SumNH4
  charge = c(0:-2, 0:-1, 0:-3, 0:-2, 0:-2, 0:-2, 0:-1, 0:-
1, 0:-1, -1:1, 1:0),
  alkfac = c(0:2, 0:1, -1:2, 0:2, 0:2, -2:0, -1:0, -1:0,
-1:0, 1:-1, 0:1),

  lumpsum = c(rep("SumCO2", 3), rep("SumBOH3",2), rep("SumH3PO4",4),
rep("SumSiOH4",3),
             rep("SumH2S",3), rep("SumH2SO4",3), rep("SumHF",2),
rep("SumHNO3",2),
             rep("SumHNO2",2), rep("SumH2O",3), rep("SumNH4",2))
)

  expression = list(expression("CO"[2]), expression("HCO"[3]^"-
"), expression("CO"[3]^"2-"), expression("BOH"[3]), expression("BOH"[4]^"-
"),
                    expression(paste("H"[3],"PO"[4])),
expression(paste("H"[2],"PO"[4]^"-")),
                    expression("HPO"[4]^"-"), expression("PO"[4]^"3-
"),
                    expression("SiOH"[4]),
expression("SiOOH"[3]^"-"), expression(paste("SiO"[2],"OH"[2]^"2-")),
                    expression(paste("H"[2],"S")), expression("HS"^"-
"), expression("S"^"2-"),
                    expression(paste("H"[2],"SO"[4])),
expression("HSO"[4]^"-"), expression("SO"[4]^"2-"),
                    expression("HF"), expression("F"^"-"),
                    expression("HNO"[3]),
expression("NO"[3]^"-"),
                    expression("HNO"[2]),
expression("NO"[2]^"-"),
                    expression("OH"^"-"),
expression(paste("H"[2],"O")), expression("H"^"+"),
                    expression("NH"[4]^"+"), expression("NH"[3]))

  Speciation$expression <- expression

  knitr::kable(Speciation[, 1:4])

...

```{r}

get.expression <- function(species = NULL, lumpsum = NULL){
 EXP <- NULL
 if (! is.null(species))
 EXP <- Speciation$expression[Speciation$species %in% species]

```

```

 if (! is.null(lumpsum))
 EXP <- c(EXP, Speciation$expression[Speciation$lumpsum %in%
lumpsum])
 unlist(EXP)

 }

...

Charge and proton balance

Several functions are created.

* *get.fraction* calculates the contribution of species to a lumpsum
* *get.charge* calculates the charge of a lumpsum.
* *get.CBA* calculates the charge balance alkalinity of a lumpsum.
* *get.PAL* calculates the proton acceptor level of a lumpsum.

```{r}
get.fraction <- function(pH = 4.5,
                        t = 18.252,
                        S = 34.617,
                        p = 0,
                        species = c("CO2", "HCO3", "CO3", "BOH3", "BOH4",
                                "H3PO4", "H2PO4", "HPO4", "PO4",
                                "SiOH4", "SiOOH3", "SiO2OH2", "H2S",
"HS", "S2min",
                                "H2SO4", "HSO4", "SO4", "HF", "F",
"HNO3", "NO3",
                                "HNO2", "NO2", "OH", "H", "H2O",
"NH4", "NH3"),
                        lumpsum = NULL,
                        SumHF = SETTINGS$SumHF, SumH2SO4 =
SETTINGS$SumH2SO4,
                        ...) {

  AE <- aquaenv(S = S, t = t, pH = pH, p = p,
              SumCO2 = 1e-6, SumNH4 = 1e-6, SumH2S = 1e-6, SumH3PO4 =
1e-6,
              SumSiOH4 = 1e-6, SumHNO3 = 1e-6, SumHNO2 = 1e-6, SumBOH3 =
1e-6,
              SumHF = SumHF, SumH2SO4 = SumH2SO4,
              ...)

  if (!is.null(species))
    species <- as.character(species)

  if (! is.null(lumpsum))
    species <- c(species, as.character(Speciation[Speciation$lumpsum %in%
lumpsum, "species"]))

  species <- unique(species)

  if (is.null(species))
    species <- c("CO2", "HCO3", "CO3", "BOH3", "BOH4", "H3PO4", "H2PO4",
                "HPO4", "PO4", "SiOH4", "SiOOH3", "SiO2OH2", "H2S", "HS",
"S2min",
                "H2SO4", "HSO4", "SO4", "HF", "F", "HNO3", "NO3", "HNO2",
"NO2",
                "OH", "H", "H2O", "NH4", "NH3")

  W <- which(species %in% c("OH", "H2O", "pH", "H", "H2SO4", "HSO4",
"SO4", "HF", "F"))

  if (length(W)) {
    Species <- species[-W]
    Water <- species[W]
  } else {

```

```

Species <- species
Water   <- NULL
}

if (length(Species))
  fraction <- as.data.frame(AE[Species])*1e6
else
  fraction <- NULL

Cwat <- 55.4939 # concentration of SUM water

if ("OH" %in% Water)
  fraction <- cbind(fraction, OH = unlist(as.data.frame(AE["OH"]))/Cwat)

if ("H2O" %in% Water)
  fraction <- cbind(fraction, H2O = unlist(1 -
as.data.frame(AE["OH"])/Cwat))

if ("H" %in% Water)
  fraction <- cbind(fraction, H = unlist(10^(-
as.data.frame(AE["pH"]))))

if ("pH" %in% Water)
  fraction <- cbind(fraction, pH = as.data.frame(AE["pH"]))

SumH2SO4 <- AE[["SumH2SO4"]]
SumHF     <- AE[["SumHF"]]
if ("H2SO4" %in% Water)
  fraction <- cbind(fraction, H2SO4 =
unlist(as.data.frame(AE["H2SO4"])/SumH2SO4))
if ("HSO4" %in% Water)
  fraction <- cbind(fraction, HSO4 =
unlist(as.data.frame(AE["HSO4"])/SumH2SO4))
if ("SO4" %in% Water)
  fraction <- cbind(fraction, SO4 =
unlist(as.data.frame(AE["SO4"])/SumH2SO4))
if ("HF" %in% Water)
  fraction <- cbind(fraction, HF =
unlist(as.data.frame(AE["HF"])/SumHF))
if ("F" %in% Water)
  fraction <- cbind(fraction, F =
unlist(as.data.frame(AE["F"])/SumHF))

if (length(species) > 1)
  fraction <- fraction[,species] # to have same ordering as input
if (is.matrix(fraction) )
  fraction <- as.data.frame(fraction)

row.names(fraction) <- NULL

return(fraction)
}
...

```{r}
bjerrum <- function(lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
 "SumH2S", "SumH2SO4", "SumHF", "SumHNO3",
 "SumHNO2", "SumH2O", "SumNH4"),
 pH = seq(from = 0, to = 14, by = 0.1), plot = TRUE,
 legend = TRUE, ...) {
 Lump <- match.arg(lumpsum, several.ok = TRUE)
 bjerrum <- get.fraction(species = NULL, lumpsum = Lump, pH = pH)

 if (plot){
 plt <- list(...)
 if (is.null(plt)) plt <- list()
 if (is.null(plt$main)) plt$main <- Lumpsum[Lump,]$expression
 if (is.null(plt$type)) plt$type <- "l"
 if (is.null(plt$lwd)) plt$lwd <- 2
 }
}

```



```

 "SumHNO3", "SumHNO2", "SumH2O",
"SumNH4")) {
 -get.charge(pH = pH, t = t, S = S, p = p, lumpsum = lumpsum)
}
...

```{r}
get.PAL <-function(pH = 4.5, t = 18.252, S = 34.617, p = 0,
                  lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
                                "SumH2S", "SumH2SO4", "SumHF",
                                "SumHNO3", "SumHNO2", "SumH2O",
"SumNH4")){
  if (length(pH) == 1){
    Spec      <- Speciation[Speciation$lumpsum %in% lumpsum, ]
    fraction <- t(get.fraction(pH = pH, t = t, S = S, p = p,
                              species = as.character(Spec$species)))
    Total.alkalinity <- Spec$alkfac * fraction
    tapply(Total.alkalinity, INDEX = Spec$lumpsum, FUN = sum)[lumpsum]
  } else {

    Alk <- NULL

    for (Lump in lumpsum){
      Spec <- Speciation[Speciation$lumpsum == Lump, ]
      fraction <- t(get.fraction(pH = pH, t = t, S = S, p = p,
                                species = as.character(Spec$species)))
      Ch <- Spec$alkfac*fraction
      CS <- colSums(Ch)

      Alk <- cbind(Alk, CS)
    }
    if (is.vector(Alk))
      Alk <- as.matrix(Alk)
    colnames(Alk) <- lumpsum
    return(Alk)
  }
}
...

# Charge and proton acceptor level as a function of pH

```{r, fig.width = 8, fig.height = 12}
Charge <- get.charge(pH = pH.seq)
Charge <- as.data.frame(Charge)
Charge$CBA <- 1
Charge$pH <- pH.seq

alkalinity <- get.PAL(pH = pH.seq)
alkalinity <- as.data.frame(alkalinity)
alkalinity$pH <- pH.seq
alkalinity$TA <- 1

P_C <- bjerrum("SumCO2", pH = pH.seq, plot = FALSE)
cnames <- c("pH", "SumCO2", "SumBOH3", "SumH3PO4", "SumNH4",
 "SumH2S", "SumH2SO4", "SumHF", "SumHNO2", "SumHNO3",
"SumSiOH4")

PAL <- alkalinity[,cnames]
CH <- Charge[,cnames]

par (mar = c(4,4,4,7), mfrow = c(3,1), las = 1)
matplot(x = pH.seq, y = P_C, type = "l", lty = 1, lwd = 2, xlab = "pH",
ylab = "-",

```

```

 main = "DIC speciation", col = "grey")
CC <- alpha.col (col = 1, alpha = 0.5)
N <- length(pH.seq)
polygon(x = c(pH.seq[1],pH.seq,pH.seq[N],pH.seq[1]),y=c(0,P_C[,1],0,0),
col=CC,border=NA)
polygon(x = c(pH.seq[1],pH.seq,pH.seq[N],pH.seq[1]),y=c(0,P_C[,2],0,0),
col=CC,border=NA)
polygon(x = c(pH.seq[1],pH.seq,pH.seq[N],pH.seq[1]),y=c(0,P_C[,3],0,0),
col=CC,border=NA)

text(3,0.97, expression(paste("H"[2],"CO"[3])))
text(7.6,0.97, expression(paste("HCO"[3]^"-")))
text(12.5,0.97, expression("CO"[3]^"2-"))

text(1.3,0.97,"(A)", cex = 1.2)

par(new = TRUE)
plot(x = PAL$pH, y = PAL$SumCO2, axes = FALSE, xlab = "", ylab="", lty = 1,
lwd = 2, type = "l")
axis (side = 4)
mtext(outer = FALSE, "Proton acceptor level", side = 4, line =-1, las=0)
par(new = TRUE) #charge
plot(x = CH$pH, y = CH$SumCO2, type = "l", lty = 2, lwd = 2, col =
"darkred", axes = FALSE, xlab="",ylab="")
axis(side=4, line = 4, col = "darkred", lty = 2, lwd = 2)
mtext(outer = FALSE, "Charge", side = 4, line =3, las=0, col = "darkred")
abline (lty = 2, v = 4.5)

LTY <- c(rep(1,4),rep(2,4),rep(3,2))
COLS <- c(2,4,8,10)

matplot(PAL$pH, PAL[,-1], type = "l", lwd = 2, lty = LTY, las = 1, col =
COLS,
 xlab = "pH", main = "Proton acceptor level", ylab = "-")
text(5.2,0.5,col=2,bty="o",expression(sum("CO"[2])))
text(8.0,0.5,col=4,expression(sum("B(OH)"[3])))
text(9,2,col=8,expression(sum(paste("H"[3], "PO"[4])))
text(14,0.9,col=10,expression(sum("NH"[4])))
text(8.8,0.9,col=2,expression(sum(paste("H"[2], "S"))))
text(1.1,-0.6,col=4,expression(sum(paste("H"[2], "SO"[4])))
text(1.7,-0.75,col=8,expression(sum("HF")))
text(3.35,-0.5,col=10,expression(sum("HNO"[2])))
text(14,-0.1,col=2,expression(sum("HNO"[3])))
text(14,1.7,col=4,expression(sum("Si(OH)"[4])))

text(1,2,"(B)", cex = 1.2)

matplot(CH$pH, CH[,-1], type = "l", lwd = 2, lty = LTY, las = 1, col =
COLS,
 xlab = "pH", main = "Charge", ylab = "-")
text(5.4,-0.5,col=2,expression(sum("CO"[2])))
text(7.8,-0.3,col=4,expression(sum("B(OH)"[3])))
text(6.7,-1.5,col=8,expression(sum(paste("H"[3], "PO"[4])))
text(13.5,0.2,col=10,expression(sum("NH"[4])))
text(7.8,-0.7,col=2,expression(sum(paste("H"[2], "S"))))
text(2.8,-1.8,col=4,expression(sum(paste("H"[2], "SO"[4])))
text(2.25,-0.5,col=8,expression(sum("HF")))
text(3.35,-0.5,col=10,expression(sum("HNO"[2])))
text(2,-1.1,col=2,expression(sum("HNO"[3])))
text(13.5,-1.5,col=4,expression(sum("Si(OH)"[4])))

text(14,1,"(C)", cex = 1.2)
...

```

```
Charge change by processes.
```

This shows the total charge of species that are involved in a certain reaction and that need to be adjusted by proton uptake/release.

It is similar as the figures in the Soetaert et al. (2007) paper, but then without the multiplication with the "sensitivity" factor.

```

```{r}
ProcessCoeff <- function(SumCO2=0, SumBOH3=0, SumH3PO4=0, SumSiOH4=0,
                        SumH2S=0, SumH2SO4=0, SumHF=0,
                        SumHNO3=0, SumHNO2=0, SumH2O=0, SumNH4=0, #
change in concentration of lumpsum species
                        TA = 0, # change in total alkalinity
                        CBA = 0) # change in charge balance
(excess negative charge)
{
  c(SumCO2 = SumCO2, SumBOH3=SumBOH3, SumH3PO4=SumH3PO4, SumSiOH4=SumSiOH4,
    SumH2S=SumH2S, SumH2SO4=SumH2SO4, SumHF=SumHF, SumHNO3=SumHNO3,
    SumHNO2=SumHNO2, SumH2O=SumH2O, SumNH4=SumNH4, TA=TA, CBA=CBA)
}

ProcessCoefficients <- t(data.frame(
  OxidationMineralisation = ProcessCoeff(SumCO2=1, SumNH4=NC, SumH3PO4=PC,
TA=NC-PC, CBA=0),
  Denitrification = ProcessCoeff(SumCO2=1, SumNH4=NC, SumH3PO4=PC,
SumHNO3=-0.8, TA=0.8+NC-PC, CBA=0),
  Denitrification2 = ProcessCoeff(SumCO2=1, SumH3PO4=PC,
SumHNO3=-(0.8+0.6*NC), TA=0.8+0.6*NC-PC, CBA=0),
  Feoxidation = ProcessCoeff(SumCO2=1, SumNH4=NC, SumH3PO4=PC,
TA=NC-PC+8, CBA=8),
  Mnoxidation = ProcessCoeff(SumCO2=1, SumNH4=NC, SumH3PO4=PC,
TA=NC-PC+4, CBA=4),
  Sulfatereduction = ProcessCoeff(SumCO2=1, SumNH4=NC, SumH3PO4=PC,
SumH2SO4=-0.5, SumH2S=0.5, TA=NC-PC+1, CBA=0),
  Methanogenesis = ProcessCoeff(SumCO2=0.5, SumNH4=NC, SumH3PO4=PC,
TA=NC-PC, CBA=0),

  Nitrification = ProcessCoeff(SumNH4=-1, SumHNO3=1, TA=-2,
CBA=0),
  Anammox = ProcessCoeff(SumNH4=-1, SumHNO2=-1, TA=0,
CBA=0),
  MnreoxidationO2 = ProcessCoeff(TA=-2,
CBA=-2),
  FereoxidationO2 = ProcessCoeff(TA=-2,
CBA=-2),
  FereoxidationNO3 = ProcessCoeff(SumHNO3=-0.2, TA=-
1.8, CBA=-2), # NOTE: WRONG IN TABLE SOETAERT ET AL!
  FereoxidationMn = ProcessCoeff(TA=-1,
CBA=-1),
  SreoxidationO2 = ProcessCoeff(SumH2S=-1, SumH2SO4=1, TA=-2,
CBA=0),
  MethaneoxidationO2 = ProcessCoeff(SumCO2=1, TA=0,
CBA=0),
  AOM = ProcessCoeff(SumH2SO4=-1, SumH2S=1, SumCO2=1, TA=2,
CBA=0),

  FeSoxidationO2 = ProcessCoeff(SumH2SO4=1, TA=-2, CBA=0),
  FeSoxidationMn = ProcessCoeff(SumH2SO4=1, TA=8, CBA=10),
  FeSoxidationFe = ProcessCoeff(TA=6, CBA=6),
  FeSprecipitation = ProcessCoeff(SumH2S=-1.5, TA=0, CBA=0),
  FeSprecipitationFe = ProcessCoeff(SumH2S=-1, TA=-2, CBA=-2),
  FeS2formation = ProcessCoeff(SumH2S=-1, TA=0, CBA=0),
  MnCO3formation = ProcessCoeff(SumCO2=-1, TA=-2, CBA=-2),
  FeCO3formation = ProcessCoeff(SumCO2=-1, TA=-2, CBA=-2),
  CaSO4formation = ProcessCoeff(SumH2SO4=-1, TA=0, CBA=-2),
  S0formationFe = ProcessCoeff(SumH2S=-1, TA=4, CBA=4),
  S0formationMn = ProcessCoeff(SumH2S=-1, TA=2, CBA=2),
  Adsorption = ProcessCoeff(TA=1, CBA=1),
  CO2release = ProcessCoeff(SumCO2=-1, TA=0, CBA=0),
  NH3release = ProcessCoeff(SumNH4=-1, TA=-1, CBA=0),
  NH4release = ProcessCoeff(SumNH4=-1, TA=0, CBA=1),

```

```

Primaryproduction = ProcessCoeff(SumCO2=-1, SumNH4=-NC, SumH3PO4=-PC,
TA=-NC+PC, CBA=0),
NO3assimilation = ProcessCoeff(SumCO2=-1, SumHNO3=-NC, SumH3PO4=-PC,
TA=NC+PC, CBA=0),
CaCO3production = ProcessCoeff(SumCO2=-1, TA=-2, CBA=-2),
CaCO3dissolution = ProcessCoeff(SumCO2=1, TA=2, CBA=2)
)
...

```{r}
dTA<- function(pH = 1:12, t = 18.252, S = 34.617, p = 0,
 SumCO2=0, SumBOH3=0, SumH3PO4=0, SumSiOH4=0,
 SumH2S=0, SumH2SO4=0, SumHF=0,
 SumHNO3=0, SumHNO2=0, SumH2O=0, SumNH4=0, # change in
concentration of lumpsum species
 TA = 0,
 process = ProcessCoeff(SumCO2, SumBOH3, SumH3PO4, SumSiOH4,
 SumH2S, SumH2SO4, SumHF,
 SumHNO3, SumHNO2, SumH2O, SumNH4, TA
= TA)
) {
 pCoeff <- process[process!=0] # Coefficients that are not 0
and not CBA
 pCoeff <- pCoeff[names(pCoeff) != "CBA"]

 lumpsum <- names(pCoeff)
 lumpsum <- lumpsum[!lumpsum %in% c("TA")]

 if (length(lumpsum))
 ALK <- -get.PAL(pH = pH, t = t, S = S, p = p, lumpsum = lumpsum) #
PAL at the pH for the relevant lump sums
 else
 ALK <- NULL

 if ("TA" %in% names(pCoeff)) ALK <- cbind(ALK, TA = 1) # if
TA is added
 if (is.matrix(ALK)){
 dTA <- t(t(ALK) * pCoeff[colnames(ALK)])
 return(rowSums(dTA))
 } else if(length(lumpsum) > 1)
 return (sum(ALK * pCoeff[names(ALK)]))
 else
 return (ALK * pCoeff)
}
...

```{r}
dCBA <- function(pH = 1:12, t = 18.252, S = 34.617, p = 0,
              SumCO2=0, SumBOH3=0, SumH3PO4=0, SumSiOH4=0,
              SumH2S=0, SumH2SO4=0, SumHF=0,
              SumHNO3=0, SumHNO2=0, SumH2O=0, SumNH4=0, # change in
concentration of lumpsum species
              CBA = 0,
              process = ProcessCoeff(SumCO2, SumBOH3, SumH3PO4,
SumSiOH4,
              SumH2S, SumH2SO4, SumHF,
              SumHNO3, SumHNO2, SumH2O, SumNH4,
CBA = CBA)
              ) {
  pCoeff <- process[process!=0]
  pCoeff <- pCoeff[names(pCoeff) != "TA"]

  lumpsum <- names(pCoeff)
  lumpsum <- lumpsum[!lumpsum %in% c("CBA")]
  if (length(lumpsum))
    CBA <- get.charge(pH = pH, t = t, S = S, p = p, lumpsum = lumpsum)
  else
    CBA <- NULL
  if ("CBA" %in% names(pCoeff)) CBA <- cbind(CBA, CBA = 1)

  if (is.matrix(CBA)){

```

```

dCBA <- t( t(CBA) * pCoeff[colnames(CBA)])
return(rowSums(dCBA))
} else if(length(lumpsum) > 1)
return (sum(CBA * pCoeff[names(CBA)]))
else
return (CBA * pCoeff)
}
...

# Bufferfactors and ENC

```{r}
dCBAdH.function <- function(ae, lumpsum = c("SumCO2", "SumBOH3",
"SumH3PO4", "SumSiOH4",
"SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",
"SumNH4"))
{
with(as.list(ae),
{
H <- 10^(-pH)

dENCdH <- (-1/H) * (OH + H) # Internal enhancement of
buffering

if ("SumCO2" %in% lumpsum & SumCO2 >0)
dENCdH <- dENCdH + (-1/H) * (HCO3 * (c1 - c3) + 2 * CO3 * (2 *
c1 + c2))

if ("SumBOH3" %in% lumpsum & SumBOH3 >0)
dENCdH <- dENCdH + (-1/H) * (BOH4 * b1)

if ("SumH3PO4" %in% lumpsum & SumH3PO4 >0)
dENCdH <- dENCdH + (-1/H) * (H2PO4 * (p1 - p3 - 2 * p4) +
2 * HPO4 * (2 * p1 + p2 - p4) + 3 * PO4 * (3 *
p1 + 2 * p2 + p3))

if ("SumSiOH4" %in% lumpsum & SumSiOH4 >0)
dENCdH <- dENCdH + (-1/H) * (SiOOH3 * (si1 -si3) + 2*SiO2OH2 *
(2*si1 + si2))

if ("SumH2S" %in% lumpsum & SumH2S > 0)
dENCdH <- dENCdH + (-1/H) * (HS * (s1 - s3) + 2 * S2min * (2 *
s1 + s2))

if ("SumH2SO4" %in% lumpsum & SumH2SO4 >0)
dENCdH <- dENCdH + (-1/H) * (HSO4 * (so1 - so3) + 2 * SO4 * (2 *
so1 + so2))

if ("SumHF" %in% lumpsum & SumHF >0)
dENCdH <- dENCdH + (-1/H) * (ae[["F"]] * f1)

if ("SumHNO3" %in% lumpsum & SumHNO3 >0)
dENCdH <- dENCdH + (-1/H) * (NO3 * na1)

if ("SumHNO2" %in% lumpsum & SumHNO2 > 0)
dENCdH <- dENCdH + (-1/H) * (NO2 * ni1)

if ("SumNH4" %in% lumpsum & SumNH4 > 0)
dENCdH <- dENCdH + (-1/H) * (-NH4 * n2)

return(dENCdH)
})
}

dHdCBA.function <- function(ae, lumpsum = c("SumCO2", "SumBOH3",
"SumH3PO4", "SumSiOH4",
"SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",
"SumNH4"))
1/dCBAdH.function (ae, lumpsum = lumpsum)

```

```

dpH.dCBA <- function(ae, lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
 "SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",
"SumNH4"))
 (-log(10)*as.numeric(10^(-ae$pH)) * dCBA.dH.function(ae = ae, lumpsum =
lumpsum))^(1)
...

```{r}
dTAdH.function <- function(ae, lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
          "SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",
"SumNH4"))
{
  with(as.list(ae),
    {
      H <- 10^(-pH)
      dTAdH <- (-1/H) * (OH + H)      # Internal enhancement of
buffering

      if ("SumCO2" %in% lumpsum & SumCO2 > 0)
        dTAdH <-dTAdH + (-1/H) * (HCO3 * (c1 - c3) + 2 * CO3 * (2 * c1 +
c2))

      if ("SumBOH3" %in% lumpsum & SumBOH3 > 0)
        dTAdH <-dTAdH + (-1/H) * (BOH4 * b1)

      if ("SumH2S" %in% lumpsum & SumH2S > 0)
        dTAdH <-dTAdH + (-1/H) * (HS * (s1 - s3) + 2 * S2min * (2 * s1 +
s2))

      if ("SumSiOH4" %in% lumpsum & SumSiOH4 > 0)
        dTAdH <-dTAdH + (-1/H) * (SiOOH3 * (si1 - si3) + 2 * SiO2OH2 *
(2 * si1 + si2))

      if ("SumNH4" %in% lumpsum & SumNH4 > 0)
        dTAdH <-dTAdH + (-1/H) * (NH3 * n1)

      if ("SumH3PO4" %in% lumpsum & SumH3PO4 > 0)
        dTAdH <-dTAdH + (-1/H) * (-H3PO4 * (-p2 - 2 * p3 - 3 * p4) +
HPO4 * (2 * p1 + p2 -
          p4) + 2 * PO4 * (3 * p1
+ 2 * p2 + p3))

      if ("SumHNO3" %in% lumpsum & SumHNO3 > 0)
        dTAdH <-dTAdH + (-1/H) * (-HNO3 * na2)

      if ("SumHNO2" %in% lumpsum & SumHNO2 > 0)
        dTAdH <-dTAdH + (-1/H) * (-HNO2 * ni2)

      if ("SumHF" %in% lumpsum & SumHF > 0)
        dTAdH <-dTAdH + (-1/H) * (-HF * f2)

      if ("SumH2SO4" %in% lumpsum & SumH2SO4 > 0)
        dTAdH <-dTAdH + (-1/H) * (-2 * H2SO4 * (-so2 - 2 * so3) - HSO4 *
(so1 - so3))

      return(dTAdH)
    })
}

dHdTA.function <- function(ae, lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
          "SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",
"SumNH4"))
  1/dTAdH.function(ae, lumpsum = lumpsum)

dpH.dTA <- function(ae, lumpsum = c("SumCO2", "SumBOH3", "SumH3PO4",
"SumSiOH4",
          "SumH2S", "SumH2SO4", "SumHF", "SumHNO3", "SumHNO2",

```

```

"SumNH4"))
  (-log(10)*as.numeric(10^(-ae$pH)) * dTAdH.function(ae = ae, lumpsum =
lumpsum))^-1)
...

```{r}
ae = with(SETTINGS, aquaenv(pH = pH.seq, SumCO2 = SumCO2, SumBOH3 =
SumBOH3,
 SumH3PO4 = SumH3PO4, SumSiOH4 = SumSiOH4, SumH2S = SumH2S,
 SumH2SO4 = SumH2SO4, SumHF = SumHF, SumHNO3 = SumHNO3,
 SumHNO2 = SumHNO2, SumNH4 = SumNH4, S = S, t = t, p = p, dsa =
TRUE))
dpHdCBA <- dpH.dCBA(ae)/1e6
dpHdTA <- dpH.dTA(ae)/1e6

species = c("SumCO2", "SumBOH3", "SumH2S", "SumSiOH4", "SumNH4",
"SumH3PO4",
 "SumHNO3", "SumHNO2", "SumHF", "SumH2SO4")
dpHdDIC <- NULL
RF <- NULL

for (pH in pH.seq) {
 ae = with(SETTINGS, aquaenv(pH = pH, SumCO2 = SumCO2, SumBOH3 = SumBOH3,
 SumH3PO4 = SumH3PO4, SumSiOH4 = SumSiOH4, SumH2S = SumH2S,
 SumH2SO4 = SumH2SO4, SumHF = SumHF, SumHNO3 = SumHNO3,
 SumHNO2 = SumHNO2, SumNH4 = SumNH4, S = S, t = t, p = p, dsa
= TRUE))
 dpHdDIC <- c(dpHdDIC,BufferFactors(ae = ae)$dpH.dtotX["SumCO2"]/1e6)
 RF <- c(RF, BufferFactors(ae = ae)$RF[1])
}
...

```{r, fig.width=8, fig.height=10}
par(mfrow = c(3,2), las = 1, mar = c(5, 5.5, 4, 0.5))

pH.peaks.CBA <-
c(pH.seq[which(dpHdCBA==max(dpHdCBA[0:(length(pH.seq)*1/3])))],

pH.seq[which(dpHdCBA==max(dpHdCBA[(length(pH.seq)*1/3+1):(length(pH.seq)*2/3])))]
,

pH.seq[which(dpHdCBA==max(dpHdCBA[(length(pH.seq)*2/3+1):length(pH.seq)]))]
)
plot(pH.seq, dpHdCBA, type = "l", lwd = 3, col = 1, xlab = "pH", ylab =
"",main = "dpH/dCBA")
mtext(expression(paste("/(",mu,"mol/kg)")),side=2, las = 0, padj = -3.4)
abline(v = pH.peaks.CBA, col = "grey")
text(14,0.0045,"(A)",cex=1.2)
plot(pH.seq, dpHdCBA, xlim = c(6,9), type = "l", lwd = 3, col = 2, xlab =
"pH", ylab = "",main = "dpH/dCBA")

pH.peaks.DIC <-
c(pH.seq[which(dpHdDIC==min(dpHdDIC[0:(length(pH.seq)*2/3])))],

pH.seq[which(dpHdDIC==min(dpHdDIC[(length(pH.seq)*2/3+1):length(pH.seq)]))]
)
plot(pH.seq, dpHdDIC, type = "l", lwd = 3, col = 1, xlab = "pH", ylab =
"",main = "dpH/dDIC")
mtext(expression(paste("/(",mu,"mol/kg)")),side=2, las = 0, padj = -3.4)
abline(v = pH.peaks.DIC, col = "grey")
text(14,-0.0034,"(B)",cex=1.2)
plot(pH.seq, dpHdDIC, xlim = c(6,9), type = "l", lwd = 3, col = 2, xlab =
"pH", ylab = "",main = "dpH/dDIC")

pH.peaks.RF <- c(pH.seq[which(RF==max(RF[0:(length(pH.seq)*2/3])))],

```

```

pH.seq[which(RF==max(RF[(length(pH.seq)*2/3+1):length(pH.seq)]))]
plot(pH.seq, RF, type = "l", lwd = 3, col = 1, xlab = "pH", ylab = "", main =
"Revelle sensitivity factor")
mtext("-", side=2, las = 0, padj = -3.4)
abline(v = pH.peaks.RF, col = "grey")
text(14,17,"(C)", cex=1.2)
plot(pH.seq, RF, xlim = c(6,9), type = "l", lwd = 3, col = 2, xlab = "pH",
ylab = "", main = "Revelle sensitivity factor")
...

```

```
## pH versus DIC/alkalinity
```

```

```{r}
DIC.seq <- seq(from = 1950, to = 2150, length.out = 100)
TA.seq <- seq(from = 2200, to = 2450, length.out = 100)
pH.mat <- outer(DIC.seq, TA.seq,
FUN = function(DIC, TA)
aquaenv(SumCO2 = DIC/1e6, SumBOH3 =SETTINGS$SumBOH3,
SumH3PO4=SETTINGS$SumH3PO4,
SumSiOH4=SETTINGS$SumSiOH4,
SumH2S=SETTINGS$SumH2S, SumH2SO4=SETTINGS$SumH2SO4,
SumHF=SETTINGS$SumHF, SumHNO3=SETTINGS$SumHNO3,
SumHNO2=SETTINGS$SumHNO2,
SumNH4=SETTINGS$SumNH4, S=SETTINGS$S,
t=SETTINGS$t, p=SETTINGS$p,
TA = TA/1e6)$pH)
pCO2.mat <- outer(DIC.seq, TA.seq,
FUN = function(DIC, TA)
aquaenv(SumCO2 = DIC/1e6, SumBOH3 =SETTINGS$SumBOH3,
SumH3PO4=SETTINGS$SumH3PO4,
SumSiOH4=SETTINGS$SumSiOH4,
SumH2S=SETTINGS$SumH2S, SumH2SO4=SETTINGS$SumH2SO4,
SumHF=SETTINGS$SumHF, SumHNO3=SETTINGS$SumHNO3,
SumHNO2=SETTINGS$SumHNO2,
SumNH4=SETTINGS$SumNH4, S=SETTINGS$S,
t=SETTINGS$t, p=SETTINGS$p,
TA = TA/1e6)$fCO2*1e6)
...

```

```

```{r, fig.width=6, fig.height=10}
require(shape)
par(mfrow = c(2,1), mar = c(4,4,2,3))
image2D(z = pH.mat, x = DIC.seq, y = TA.seq, las = 1, xlab =
expression(paste("DIC(", mu, "M)")),
clab = expression("pH"), colkey = list(length = 0.5, width = 0.5,
dist = 0.02),
ylab = expression(paste("TA(", mu, "M)")), contour = list(col =
"darkblue"),
col = ColScheme)
text(2140,2440,"(A)")

center <- c(2100, 2250)

dConc <- 30
Hplus <- center+dConc*c(0, -1)
CO2 <- center+dConc*c(1, 0)
f1 <- uniroot(f = function(x) sqrt(2*x^2)-1, c(0,dConc))$root
HCO3 <- center+dConc*c(1, 1)*f1
f2 <- uniroot(f = function(x) sqrt(1.5*x^2)-1, c(0,dConc))$root
CO3 <- center+dConc*c(0.5, 1)*f2

Arrows(center[1], center[2], Hplus[1], Hplus[2], arr.type = "triangle")
cat("pH change by proton, from", p1 <- pH.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat(" to ", p2 <- pH.mat[which.min(abs(DIC.seq - Hplus[1])),
which.min(abs(TA.seq - Hplus[2]))], " diff = ", p2-p1, "\n")

```

```

Arrows(center[1], center[2], CO3[1], CO3[2], arr.type = "triangle")
cat("pH change by CO3, from", p1 <- pH.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat(" to ", p2 <- pH.mat[which.min(abs(DIC.seq - CO3[1]))],
which.min(abs(TA.seq - CO3[2]))], " diff = ", p2-p1, "\n")

center <- c(1970, 2400)
dConc <- 30
Hplus <- center+dConc*c(0, -1)
CO2 <- center+dConc*c(1, 0)
HCO3 <- center+dConc*c(1, 1)*f1
CO3 <- center+dConc*c(0.5, 1)*f2

Arrows(center[1], center[2], Hplus[1], Hplus[2], arr.type = "triangle")
cat("pH change by proton, from", p1<-pH.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat (" to ", p2<-pH.mat[which.min(abs(DIC.seq - Hplus[1]))],
which.min(abs(TA.seq - Hplus[2]))], " diff = ", p2-p1, "\n")

Arrows(center[1], center[2], CO3[1], CO3[2], arr.type = "triangle")
cat("pH change by CO3, from", p1<-pH.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat(" to", p2<-pH.mat[which.min(abs(DIC.seq - CO3[1]))],
which.min(abs(TA.seq - CO3[2]))], " diff = ", p2-p1, "\n")

image2D(z = pCO2.mat, x = DIC.seq, y = TA.seq, las = 1, xlab =
expression(paste("DIC(",mu,"M)")),
clab = expression("pCO"[2]), colkey = list(length = 0.5, width =
0.5, dist = 0.02),
ylab = expression(paste("TA(",mu,"M)")), resfac = 4, contour =
list(col = "darkblue"),
col = ColScheme)
text(2140,2440,"(B)")

center <- c(2100, 2250)
dConc <- 30
Hplus <- center+dConc*c(0, -1)
CO2 <- center+dConc*c(1, 0)
HCO3 <- center+dConc*c(1, 1)*f1
CO3 <- center+dConc*c(0.5, 1)*f2

Arrows(center[1], center[2], Hplus[1], Hplus[2], arr.type = "triangle")
cat("pCO2 change by proton from", p1 <-pCO2.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat(" to ", p2<-pCO2.mat[which.min(abs(DIC.seq - Hplus[1]))],
which.min(abs(TA.seq - Hplus[2]))], " diff = ", p2-p1, "\n")

Arrows(center[1], center[2], CO3[1], CO3[2], arr.type = "triangle")
cat("pCO2 change in CO3, from", p1<-pCO2.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat("to ",p2<-pCO2.mat[which.min(abs(DIC.seq - CO3[1]))],
which.min(abs(TA.seq - CO3[2]))], " diff = ", p2-p1, "\n")

center <- c(1970, 2400)
dConc <- 30
Hplus <- center+dConc*c(0, -1)
CO2 <- center+dConc*c(1, 0)
HCO3 <- center+dConc*c(1, 1)*f1
CO3 <- center+dConc*c(0.5, 1)*f2

Arrows(center[1], center[2], Hplus[1], Hplus[2], arr.type = "triangle")
cat("pCO2 change by proton, from", p1<-pCO2.mat[which.min(abs(DIC.seq -
center[1]))], which.min(abs(TA.seq - center[2]))]) # corr pH value
cat("to", p2<-pCO2.mat[which.min(abs(DIC.seq - Hplus[1]))],
which.min(abs(TA.seq - Hplus[2]))], " diff = ", p2-p1, "\n")

Arrows(center[1], center[2], CO3[1], CO3[2], arr.type = "triangle")
cat("pCO2 change in CO3, from", p1<-pCO2.mat[which.min(abs(DIC.seq -

```

```

center[1])), which.min(abs(TA.seq - center[2]))] # corr pH value
cat("to", p2<-pCO2.mat[which.min(abs(DIC.seq - CO3[1])),
which.min(abs(TA.seq - CO3[2]))], " diff = ", p2-p1, "\n")
...

# Effect of biogeochemical processes on pH
```{r}

function to estimate the pH change
SETTINGS = default speciation as in Hagens

dPH.numeric <- function(Species = SETTINGS,
 dSumCO2 = 0, dSumBOH3 = 0, dSumH2S = 0, #
change in summed conc due to process
 dSumSiOH4 = 0, dSumNH4 = 0, dSumH3PO4 = 0,
default = 0
 dSumHNO3 = 0, dSumHNO2 = 0, dSumHF = 0, dSumH2SO4
= 0,
 dTA = 0, dC = 1e-8 ,
change in total alkalinity due to process
 pHSEQ = pH.seq) { # pH range
for which to estimate dpHdProcess

unless otherwise mentioned the change in species composition = 0
dS <- list(SumCO2 = dSumCO2, SumBOH3 = dSumBOH3, SumH3PO4 =
dSumH3PO4, # change in summed conc due to process
 SumSiOH4 = dSumSiOH4, SumH2S = dSumH2S, SumH2SO4 = dSumH2SO4,
 SumHF = dSumHF,
 SumHNO3 = dSumHNO3, SumHNO2 = dSumHNO2, SumNH4 = dSumNH4,
dS = 0, dt = 0, dp = 0)

SPEC <- Species[-which(names(Species) == "SumH2O")]
SP2 <- unlist(SPEC) + dC*unlist(dS) # new concentrations, including the
small perturbation due to the process

dPH_dSpec <- NULL # will contain the ultimate results

for (pH in pHSEQ){

given the pH and the default species composition (SPEC), calculate TA
TA <- do.call("aquaenv" , c(as.list(SPEC), pH = pH))$TA

given new TA (+ the production by process) and the new species
composition (SP2), calculate pH
pH2 <- do.call("aquaenv" , c(as.list(SP2), TA = TA + dTA*dC))$pH

estimate dpHdspec by numerical differencing. Divide by 1e6 so that pH
change is per micromol rather than per mol.
dPH_dSpec <- c(dPH_dSpec, (pH2-pH)/(dC*1e6)) # units in per umolC
}
return(dPH_dSpec)
}
DENIT <- dPH.numeric(dSumCO2 = 1, dSumNH4 = NC, dSumH3PO4 = PC, dSumHNO3 =
-0.8, dTA = 0.8 + NC-PC)
...

```{r}

Numerical <- cbind(
  oxicmineralisation = dPH.numeric(dSumCO2=1, dSumNH4=NC, dSumH3PO4=PC,
dTA=NC-PC),
  denitrification = dPH.numeric(dSumCO2=1, dSumNH4=NC, dSumH3PO4=PC,
dSumHNO3=-0.8, dTA=0.8+NC-PC),
  denitrification2 = dPH.numeric(dSumCO2=1, dSumH3PO4=PC,
dSumHNO3=-(0.8+0.6*NC), dTA=0.8+0.6*NC-PC),

```

```

Feoxidation      = dPH.numeric(dSumCO2=1,    dSumNH4=NC,  dSumH3PO4=PC,
dTA=NC-PC+8),
Mnoxidation      = dPH.numeric(dSumCO2=1,    dSumNH4=NC,  dSumH3PO4=PC,
dTA=NC-PC+4),
Sulfatereduction = dPH.numeric(dSumCO2=1,    dSumNH4=NC,  dSumH3PO4=PC,
dSumH2SO4=-0.5, dSumH2S=0.5, dTA=NC-PC+1),
Methanogenesis   = dPH.numeric(dSumCO2=0.5,  dSumNH4=NC,  dSumH3PO4=PC,
dTA=NC-PC),

Nitrification    = dPH.numeric(dSumNH4=-1,  dSumHNO3=1,          dTA=-
2),
Anammox          = dPH.numeric(dSumNH4=-1,  dSumHNO2=-1,        dTA=0),
Mnreoxidation    = dPH.numeric(              dTA=-
2),
Fereoxidation    = dPH.numeric(              dTA=-
2),
FereoxidationNO3 = dPH.numeric(dSumHNO3=-0.2, dTA=-
1.8), # NOTE: WRONG IN TABLE SOETAERT ET AL!
FereoxidationMn  = dPH.numeric(              dTA=-
1),
Sreoxidation     = dPH.numeric(dSumH2S=-1,  dSumH2SO4=1,        dTA=-
2),
Methaneoxidation = dPH.numeric(dSumCO2=1,
dTA=0),
AOM              = dPH.numeric(dSumH2SO4=-1, dSumH2S=1,  dSumCO2=1,
dTA=2),

FeSoxidation     = dPH.numeric(dSumH2SO4=1,  dTA=-2),
FeSoxidationMn   = dPH.numeric(dSumH2SO4=1,  dTA=8),
FeSoxidationFe   = dPH.numeric(              dTA=6),
FeSprecipitation = dPH.numeric(dSumH2S=-1.5, dTA=0),
FeSprecipitationFe = dPH.numeric(dSumH2S=-1,  dTA=-2),
FeS2formation    = dPH.numeric(dSumH2S=-1,  dTA=0),
MnCO3formation   = dPH.numeric(dSumCO2=-1,  dTA=-2),
FeCO3formation   = dPH.numeric(dSumCO2=-1,  dTA=-2),
CaSO4formation   = dPH.numeric(dSumH2SO4=-1, dTA=0),
S0formationFe    = dPH.numeric(dSumH2S=-1,  dTA=4),
S0formationMn    = dPH.numeric(dSumH2S=-1,  dTA=2),
adsorption       = dPH.numeric(              dTA=1),
CO2release       = dPH.numeric(dSumCO2=-1,  dTA=0),
NH3release       = dPH.numeric(dSumNH4=-1,  dTA=-1),
NH4release       = dPH.numeric(dSumNH4=-1,  dTA=0),

primaryproduction = dPH.numeric(dSumCO2=-1,  dSumNH4=-NC,  dSumH3PO4=-PC,
dTA=-NC+PC),
NO3assimilation  = dPH.numeric(dSumCO2=-1,  dSumHNO3=-NC,  dSumH3PO4=-PC,
dTA=NC+PC),
CaCO3production  = dPH.numeric(dSumCO2=-1,  dTA=-2),
CaCO3dissolution = dPH.numeric(dSumCO2=1,    dTA=2)
)

OXIC  <- dPH.numeric(dSumCO2 = 1, dSumNH4 = NC, dSumH3PO4 = PC, dTA = NC-
PC)
OXIC2 <- dPH.numeric(dSumCO2 = 1, dSumHNO3= NC, dSumH3PO4 = PC, dTA = -NC-
PC)
SULF  <- dPH.numeric(dSumCO2 = 1, dSumNH4 = NC, dSumH3PO4 = PC, dTA = 1 +
NC-PC)
METH  <- dPH.numeric(dSumCO2 =0.5, dSumNH4 = NC, dSumH3PO4 = PC, dTA =
NC-PC)

PPROD <- dPH.numeric(dSumCO2 = -1, dSumNH4 =-NC, dSumH3PO4 =-PC, dTA = -
NC+PC)
PPROD2<- dPH.numeric(dSumCO2 = -1, dSumHNO3=-NC, dSumH3PO4 =-PC, dTA =
NC+PC)

CALC  <- dPH.numeric(dSumCO2 = -1, , dTA = -2)
DISS  <- dPH.numeric(dSumCO2 = 1, , dTA = 2)
ANNAMOX <- dPH.numeric(dSumNH4 = -1, dSumHNO2 = -1, dTA = 0)

```

```
# Point of calcification/dissolution switch
```

```

OMEGA <- NULL
SPEC <- SETTINGS[-which(names(SETTINGS) == "SumH2O")]
for (pH in pH.seq){
  OMEGA <- c(OMEGA, do.call("aquaenv" , c(as.list(SPEC), pH =
pH))$omega_calcite)
}

crit <- which.min(abs(OMEGA-1))
DISS[crit:length(pH.seq)] <- NA
CALC[1:crit] <- NA
```



```

```{r, fig.width=8, fig.height=3}
par(mfrow = c(1,3), las = 1, mar = c(5,6,4,0))
DENIT <- dPH.numeric(dSumCO2 = 1, dSumNH4 = NC, dSumH3PO4 = PC, dSumHNO3 =
-0.8, dTA = 0.8+ NC-PC)

plot(pH.seq, dpHdCBA, type = "l", ylab = "", main = "sensitivity", xlab =
"pH")
mtext(expression(paste("/(",mu,"mol/kg)")),side=2, las = 0, padj = -
3.8,cex=0.9)
text(13.7,0.00435,"(A)")

plot(pH.seq, dCBA(pH.seq, process = ProcessCoefficients["Denitrification",
]), type = "l", ylab = "", main = expression(paste(Delta,"charge",sep="")),
xlab = "pH")
mtext("-",side=2, las = 0, padj = -3,cex=0.9)
text(1.3,-1.2,"(B)")
abline(h = 0, lty = 2)

plot(pH.seq, DENIT, type = "l", ylab = "", main = "response" , xlab = "pH")
mtext("dpH",side=2, las = 0, padj = -3.9,cex=0.9)
text(13.7,0.004,"(C)")
abline(h = 0, lty = 2)
```



```

```{r, fig.width=10, fig.height=8}
par(mfrow = c(2,2), las = 1)
ylab <- ""
ylab = "dpH"
xlab <- "pH"
col = c("black","darkgrey")
matplot(x = pH.seq, y = cbind(OXIC, OXIC2), type = "l", ylab = ylab,
xlab = xlab, lty = 1:2, lwd = 3, col = col)
abline(h=0, lty = 3)
legend("bottomleft", col = col, lty = 1:2, lwd = 2,
legend = c("Aerobic", "Aerobic+Nitrif"), cex = 0.8)
text(14, (min(cbind(OXIC, OXIC2))),"(A)")
matplot(x = pH.seq, y = cbind(SULF, METH), type = "l", ylab = ylab,
xlab = xlab, lty = 1:2, lwd = 3, col = col)
abline(h=0, lty = 3)
legend("topright", col = col, lty = 1:2, lwd = 2,
legend = c("Sulphate reduction", "Methanogenesis"), cex = 0.8)
text(14, (min(cbind(SULF, METH))),"(B)")
matplot(x = pH.seq, y = cbind(PPROD, PPROD2), type = "l", ylab = ylab,
xlab = xlab, lty = 1:2, lwd = 3, col = col)
abline(h=0, lty = 3)
legend("topleft", col = col, lty = 1:2, lwd = 2,
legend = c("Prim prod (ammonium)", "Prim prod (nitrate)"), cex =
0.8)
text(14, (min(cbind(PPROD, PPROD2))),"(C)")
matplot(x = pH.seq, y = cbind(CALC, DISS), type = "l", ylab = ylab,
xlab = xlab, lty = 1:2, lwd = 3, col = col)
abline(h=0, lty = 3)
legend("topright", col = col, lty = 1:2, lwd = 2,
legend = c("Calcification", "Dissolution"), cex = 0.8)
text(14, (min(cbind(CALC, DISS),na.rm=TRUE))),"(D)")
```

```


```


```

#### # References

Andreas F. Hofmann, Karline Soetaert, Jack J. Middelburg and Filip J. R. Meysman, 2010. AquaEnv - an Aquatic acid-base modelling Environment in R. Aquatic Geochemistry, DOI 10.1007/s10498-009-9084-1.

Middelburg, J.J., K. Soetaert and M. Hagens, 2020. Ocean alkalinity, buffering and biogeochemical processes. Reviews of Geophysics.

Hagens M. and J.J. Middelburg, 2016 Generalised expressions for the response of pH to changes in ocean chemistry. Geochimica et Cosmochimica Acta 187: 334-349. DOI: 10.1016/j.gca.2016.04.012.

Soetaert, K., AF.Hofmann, J.J.Middelburg, F.J.R.Meysman, J. Greenwood, 2007. The effect of biogeochemical processes on pH. Marine Chemistry 105, 30-51